



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N	A2	(11) International Publication Number: WO 98/00695													
		(43) International Publication Date: 8 January 1998 (08.01.98)													
<p>(21) International Application Number: PCT/US97/11713</p> <p>(22) International Filing Date: 3 July 1997 (03.07.97)</p> <p>(30) Priority Data: 60/021,218 3 July 1996 (03.07.96) US 60/033,152 13 December 1996 (13.12.96) US</p> <p>(71) Applicant (for all designated States except US): MASSACHUSETTS INSTITUTE OF TECHNOLOGY (US/US); 77 Massachusetts Avenue, Cambridge, MA 02139 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SHARP, Phillip, A. (US/US); Building E17-529, 77 Massachusetts Avenue, Cambridge, MA 02139 (US). ZHOU, Qiang (CN/US); Building E17-529, 77 Massachusetts Avenue, Cambridge, MA 02139 (US).</p> <p>(74) Agent: GATES, Edward, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).</p>		<p>(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>													
<p>(54) Title: TAT-SF: COFACTOR FOR STIMULATION OF TRANSCRIPTIONAL ELONGATION BY HIV-1 TAT</p> <div data-bbox="568 1155 1023 1617"> <table border="1"> <caption>Fold Enhancement Data</caption> <thead> <tr> <th>Factor</th> <th>Condition</th> <th>Fold Enhancement</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Tat</td> <td>VECTOR</td> <td>~1.0</td> </tr> <tr> <td>Tat-SF1</td> <td>~5.2</td> </tr> <tr> <td rowspan="2">Vp16</td> <td>VECTOR</td> <td>~1.0</td> </tr> <tr> <td>Tat-SF1</td> <td>~0.8</td> </tr> </tbody> </table> </div> <p>(57) Abstract</p> <p>The invention pertains to a novel transcriptional activity factor, Tat-Stimulatory Factor, as well as genes encoding this factor and fragments and biologically functional variants thereof. The Tat-Stimulatory Factor is involved in the regulation of transcriptional elongation of HIV-1 by Tat. The invention also pertains to therapeutics involving the foregoing proteins and genes, and agents that bind to the foregoing proteins and genes. The invention also relates to methods of screening for a compound which binds to Tat-SF1, Tat-SF1-associated kinase and/or a complex of Tat-SF1 and Tat-SF1-associated kinase, as well as methods of screening for compounds which modulate Tat-SF1-mediated transcriptional activation.</p>			Factor	Condition	Fold Enhancement	Tat	VECTOR	~1.0	Tat-SF1	~5.2	Vp16	VECTOR	~1.0	Tat-SF1	~0.8
Factor	Condition	Fold Enhancement													
Tat	VECTOR	~1.0													
	Tat-SF1	~5.2													
Vp16	VECTOR	~1.0													
	Tat-SF1	~0.8													

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**TAT-SF: COFACTOR FOR STIMULATION OF
TRANSCRIPTIONAL ELONGATION BY HIV-1 TAT**

Government Support

This work was funded in part by the National Institutes of Health under the Grant Nos. GM34277 and AI32486, and the National Cancer Institute under Center core Grant No. CA14051. The Government may retain certain rights in this invention.

Background of the Invention

Intricate mechanisms regulate mRNA synthesis by control of initiation or elongation of transcription. An understanding of these mechanisms and the factors controlling these mechanisms would be important in designing therapeutic modalities for treating a variety of important medical conditions, including cancer and infection. For example, HIV-1 transcriptional elongation by Tat is essential for viral replication. Interruption of transcriptional elongation by Tat, therefore, would be highly desirable as a means for treating HIV-infected individuals.

Tat activation of HIV-1 transcription is mechanistically different from conventional activation of transcription by DNA sequence-specific transcription factors. First, most conventional activators affect transcription primarily through increasing the rate of initiation, although recent studies indicate that some prototype DNA sequence-specific transcription factors such as GAL4-VP16 can stimulate both initiation and elongation. In contrast, Tat predominantly stimulates the efficiency of elongation. Secondly, while most conventional activators interact with promoter or enhancer DNA, Tat interacts with the *trans*-acting responsive (TAR) RNA element. TAR is located at the 5' end of the nascent viral transcript and forms a stem-loop structure. The specific binding of Tat to TAR depends primarily upon the integrity of the bulge loop and immediately flanking sequences in the double-stranded RNA. Sequences in the apical loop of TAR are also important for Tat activation of transcription *in vivo*.

Control of transcriptional elongation thus has been recognized as an important step in gene regulation, but mechanisms regulating the efficiency of elongation, mediated by RNA polymerase II, have not been extensively studied. The necessity for strict control of elongation for proper gene regulation is further highlighted by the recent finding that an elongation factor, Elongin, is probably the functional target of the von Hippel-Lindau tumor suppressor protein.

Summary of the Invention

The invention involves in one respect the identification, purification, and isolation of proteins, Tat-Stimulatory Factor protein, which are specifically required for Tat *trans*-activation. The invention also involves nucleic acid molecules encoding those proteins. The invention
5 further involves the discovery and identification of kinases that bind the Tat-Stimulatory Factor proteins, which binding is believed important for TAT transcriptional elongation. The expression and biological activity of the proteins are necessary for transcriptional elongation, and alteration of the expression or biological activity of these proteins can be used to influence transcriptional activity, and thereby affect critical cellular processes.

10 The preferred nucleic acids of the invention are homologues and alleles of the coding region of the nucleic acid of SEQ ID NO:1. The invention further embraces functional equivalents, variants, analogues and fragments of the foregoing nucleic acids and also embraces proteins and peptides coded for by any of the foregoing.

According to one particular aspect of the invention, an isolated nucleic acid molecule is
15 provided. The molecule hybridizes under stringent conditions to a molecule consisting of the coding region of the nucleic acid sequence of SEQ ID NO:1 and it codes for a Tat-Stimulatory Factor protein. The invention further embraces nucleic acid molecules that differ from the foregoing isolated nucleic acid molecules in codon sequence due to the degeneracy of the genetic code. The invention also embraces complements of the foregoing nucleic acids. Preferred
20 isolated nucleic acid molecules are those comprising the human cDNAs or genes corresponding to SEQ ID NO:1. Unique fragments of the foregoing molecules are specifically contemplated by the inventors.

The invention in another aspect involves expression vectors, and host cells transformed or transfected with such expression vectors, comprising the nucleic acid molecules described above.
25 In one embodiment of the invention, the host cell is a hematopoietic T-cell precursor, such as a stem cell, and the nucleic acid is an antisense nucleic acid or a nucleic acid encoding a dominant negative mutant of the Tat-Stimulating Factor protein.

According to another aspect of the invention, an isolated nucleic acid molecule is provided which comprises a unique fragment of SEQ ID NO:1. In one embodiment the unique
30 fragment is a portion of the segment of SEQ ID NO:1 consisting of SEQ ID NO:3. In another embodiment it is a portion of the segment of SEQ ID NO:1 beginning at nucleotide number 53 and ending at nucleotide number 2703, wherein the fragment is between 12 and 2650 nucleotides

in length, and complements thereof. In one embodiment, the unique fragment is at least 150 and, more preferably, at least 200 nucleotides in length. In another embodiment, the unique fragment is between 12 and 32 contiguous nucleotides in length. In all embodiments the unique fragment includes consecutive nucleotides of SEQ ID NO:1 other than the nucleotides of SEQ ID NO:1

5 which code for SEQ ID NO:3.

According to another aspect of the invention, isolated polypeptides coded for by the isolated nucleic acid molecules described above also are provided as well as functional equivalents, variants, analogs and fragments thereof. In one embodiment, the polypeptide is a human Tat-Stimulatory Factor protein or a functionally active fragment or variant thereof. In
10 another embodiment the polypeptide is a dominant negative mutant of a Tat-Stimulatory Factor protein.

The invention in another aspect involves a method for influencing transcription in a cell. An agent that selectively binds to an isolated nucleic acid molecule as described above or an expression product thereof is introduced within a cell, in an amount effective to alter
15 transcription in the cell. Preferred agents are modified antisense nucleic acids and polypeptides. In one embodiment, transcriptional elongation activity altered, and in one particularly important embodiment, HIV-1 transcriptional elongation by Tat is altered. In this embodiment, the transcriptional elongation activity can be altered to treat an individual who is infected by HIV.

The invention in another aspect involves a method for isolating a kinase. A solution
20 suspected of containing the kinase is contacted with a Tat-Stimulatory Factor protein or functional fragment thereof, and a material that binds to the Tat-Stimulatory Factor protein and that has kinase activity is identified and isolated.

The invention in a related aspect involves isolated kinases that are the binding partners of Tat-Stimulatory Factor proteins, and nucleic acids which encode such kinases, as well as
25 functional fragments, variants and analogs of the foregoing.

The invention also provides isolated polypeptides which selectively bind a Tat-Stimulatory Factor protein, a kinase binding partner of a Tat-Stimulatory Factor protein or fragments thereof. Isolated binding polypeptides include antibodies and fragments of antibodies (e.g. Fab, F(ab)₂, Fd and antibody fragments which include a CDR3 region which binds
30 selectively to a Tat-Stimulatory Factor protein or fragment thereof).

The invention also contemplates gene therapy for HIV-infected individuals, wherein stem cells of a donor are genetically engineered to include an agent that selectively binds to a nucleic

acid molecule encoding a Tat-Stimulatory Factor protein or an expression product thereof, whereby said recombinant stem cells are resistant to intracellular HIV replication.

The invention involves methods of screening for compounds which bind to Tat-SF1, Tat-SF1 associated kinase, or a complex of Tat-SF1 and its associated kinase. The invention also involves methods for screening for compounds which modulate Tat-SF1 - dependent transcriptional activation. Compounds identified by the methods of the invention are useful for detecting the presence of and/or modulating the activity of Tat-SF1, Tat-SF1 associated kinase, or a complex of Tat-SF1 and its associated kinase.

According to one aspect of the invention, a method of screening for a compound which binds to Tat-SF1 is provided. Tat-SF1 is contacted with the compound and the binding of the compound to Tat-SF1 is determined. The compound can be detectably labeled. In this preferred embodiment, determining the binding involves detecting the labeled compound bound to Tat-SF1. In another embodiment, determination of the binding of the compound to Tat-SF1 includes detecting a change in the biological activity of Tat-SF1. Preferably, Tat-SF1 biological activity is assayed by a Tat-SF1 mediated transcription assay, a Tat-SF1 immunoassay and/or a Tat-SF1-TAR binding assay in the presence and absence of the compound. In certain of the foregoing embodiments, the compound being screened which binds to Tat-SF1 is an oligonucleotide.

According to another aspect of the invention, a method for screening compounds which bind to Tat-SF1 associated kinase is provided. The Tat-SF1 associated kinase is contacted with the compound and the binding of the compound to the Tat-SF1 associated kinase is determined. In one embodiment, the compound is detectably labeled, and determining binding involves detecting the labeled compound bound to Tat-SF1. In another embodiment, determining the binding of the compound to the Tat-SF1 associated kinase involves detecting a change in the biological activity of the Tat-SF1 associated kinase. Preferably, the change in the biological activity of the Tat-SF1 associated kinase is determined by a Tat-SF1 mediated transcription assay, a Tat-SF1 associated kinase immunoassay, or a Tat-SF1 associated kinase substrate phosphorylation assay. In certain of the foregoing embodiments, the compound being screened which binds to the Tat-SF1 associated kinase is an oligonucleotide.

According to still another aspect of the invention, a method for screening a compound which binds to a complex of Tat-SF1 and Tat-SF1 associated kinase is provided. A complex of Tat-SF1 and Tat-SF1 associated kinase is contacted with the compound and the binding of the compound to the complex is determined. The compound can be detectably labeled, and

determining binding involves detecting the labeled compound bound to Tat-SF1. In another embodiment, determining the binding involves detecting a change in the biological activity of the complex of Tat-SF1 and Tat-SF1 associated kinase. Preferably, a change in the biological activity of the complex is determined by a Tat-SF1 mediated transcription assay, an
5 immunoassay of the complex of Tat-SF1 and Tat-SF1 associated kinase, a Tat-SF1 associated kinase substrate phosphorylation assay, or a Tat-SF1-TAR binding assay. In certain of the foregoing embodiments, the compound being screened which binds to the complex of Tat-SF1 and Tat-SF1 associated kinase is an oligonucleotide.

According to still another aspect of the invention, a method for screening compounds
10 which modulate Tat-SF1 dependent transcriptional activation is provided. A mammalian cell which includes a gene encoding a Tat-SF1 polypeptide is provided and contacted with the compound. Tat-SF1 mediated transcriptional activation is determined as a measure of the modulation in Tat-SF1 mediated transcription caused by contact with the compound. Preferably, the mammalian cell also includes a gene encoding a TAT polypeptide. More preferably, the
15 mammalian cell also includes an indicator gene encoding an indicator gene product operably linked to a TAR element. Still more preferably, the mammalian cell includes a gene encoding a Tat-SF1 associated kinase polypeptide. In certain embodiments, any one or more of or all of the Tat-SF1 polypeptide, the TAT polypeptide, the indicator gene product and/or the Tat-SF1 associated kinase polypeptide are encoded by transfected expression vectors.

20 In certain embodiments, the indicator gene encodes beta-galactosidase, alkaline phosphatase, chloramphenicol acetyl transferase, luciferase, or green fluorescent protein.

In certain preferred embodiments, the TAR element operably linked to the indicator gene is an HIV-1 LTR TAR element.

These and other objects and features of the invention are described in greater detail
25 below.

Brief Description of the Figures

Fig. 1 shows the identification of Tat-SF activity in cellular extracts. A. Tat activation of HIV transcriptional elongation requires a cellular activity, Tat-SF. B. Detection of the
30 phosphorylated pp140 on an immobilized HIV-1 TAR RNA. C. The cysteine-rich activation domain of Tat is required for pp140 phosphorylation on TAR.

Fig. 2 shows that Tat-SF transcriptional activity and pp140 co-peaked during glycerol gradient sedimentation. A. Detection by transcription reaction. B. Detection by kinase assay.

Fig 3 depicts purification of pp140 and Tat-SF transcriptional activity by Tat affinity columns. A. Eluates of the columns were tested for transcription activity. B. The same eluates
5 were tested for the presence of pp140 in a kinase assay. C. A silver-stained SDS gel of the above two fractions is shown.

Fig. 4 shows that the presence of pp140 is required for Tat-SF activity. A. pp140 and a cellular kinase form a complex independently of Tat. B. Immunodepletion of pp140 from a partially purified Tat-SF fraction inactivated Tat-SF transcriptional activity. C. Anti-pp140
10 antibody efficiently removed pp140 from the Tat-SF fraction.

Fig. 5. A. Amino acid sequence and domain structure of Tat-SF1. B. Similarity between Tat-SF1 and human EWS.

Fig. 6 shows that overexpression of Tat-SF1 enhances Tat activation in HeLa cells.

15 Brief Description of Sequences

SEQ ID NO:1 is a nucleic acid including the coding region of Tat-Stimulatory Factor.

SEQ ID NO:2 is the translated amino acid sequence of the coding region of SEQ ID NO:1.

SEQ ID NO:3 is an expressed sequence tag, an amino acid sequence encoded by a portion
20 of SEQ ID NO:1.

SEQ ID NO:4 is a portion of the amino acid sequence of EWS, which shows some homology with the amino acid of SEQ ID NO:2.

SEQ ID NO:5 is a portion of the nucleic acid of SEQ ID NO:1.

25 Detailed Description of the Invention

Using a reconstituted reaction that supports a TAR-dependent and Tat-specific activation of elongation, we have identified, purified, and isolated a cDNA for a novel cellular activity, Tat-Stimulatory Factor, that is specifically required for Tat *trans*-activation. This factor is a substrate of an associated cellular kinase. Co-transfection with the cDNA for Tat-Stimulatory Factor
30 specifically stimulates Tat activation of HIV transcription. Sequence analysis indicates that Tat-Stimulatory Factor is related to EWS and FUS/TLS, which are members of a novel family of putative transcription factors with RNA recognition motifs and are frequently associated with

many types of sarcomas. It is believed that Tat activates the processivity of elongation by recruitment of a pre-formed complex containing Tat-Stimulatory Factor and a kinase to the HIV-1 promoter through a Tat-TAR interaction.

The mRNA transcript is about 3.0 kb in length, with an open reading frame of 2271 bp.

5 The open reading frame encodes protein of 754 amino acids with a calculated molecular weight of 85,767 Daltons. Sequence analysis of the protein reveals that it has several unique features. The protein can be roughly divided at position 420 into two halves. The COOH-terminal half is extremely rich in acidic amino acids, with 48% of the last 245 amino acid residues as glutamate or aspartate. The COOH-terminal half also contained many serine residues that are contained in
10 a short peptide sequence matching consensus sites for phosphorylation by Casein Kinase II. Such phosphorylation would contribute more negative charges to this region. The NH₂ terminal half of Tat-Stimulatory Factor contains two tandem RNA recognition motifs, which have homology to many RNA-binding proteins. Further details about the protein are described in greater detail in the examples below.

15 It was determined that overexpression of Tat-Stimulatory Factor enhances Tat activation *in vivo*. Immunodepletion of the Tat-Stimulatory Factor from a partially purified fraction containing Tat-Stimulatory Factor transcriptional activity eliminates its ability to support Tat *trans*-activation. Thus, it is believed that Tat-Stimulatory Factor is required for Tat *trans*-activation.

20 The invention thus involves in one aspect Tat-Stimulatory Factor proteins, genes encoding those proteins, functional modifications and variants of the foregoing, useful fragments of the foregoing, as well as therapeutics relating thereto.

Homologs and alleles of the Tat-Stimulatory Factor nucleic acids of the invention can be identified by conventional techniques. Thus, an aspect of the invention is those nucleic acid
25 sequences which code for Tat-Stimulatory Factor proteins and which hybridize to a nucleic acid molecule consisting of the coding region of SEQ ID NO:1, under stringent conditions. The term "stringent conditions" as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold
30 Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. More specifically, stringent condition, as used herein, refers to hybridization at 65°C in hybridization

buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% Bovine Serum Albumin, 2.5mM NaH₂PO₄(pH7), 0.5% SDS, 2mM EDTA). SSC is 0.15M sodium chloride/0.15M sodium citrate, pH7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid. After hybridization, the membrane upon which the DNA is transferred is washed at 2 x
5 SSC at room temperature and then at 0.1 x SSC/0.1 x SDS at 65°C.

There are other conditions, reagents, and so forth which can be used, which result in a similar degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to manipulate the conditions in a manner to permit the clear identification of homologs and alleles
10 of Tat-Stimulatory Factor nucleic acids of the invention. The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules which then are routinely isolated, followed by isolation of the pertinent nucleic acid molecule and sequencing.

In general homologs and alleles typically will share at least 40% nucleotide identity
15 and/or at least 50% amino acid identity to the coding region of SEQ ID NOs:1 or 2 (Figure 5), respectively, in some instances will share at least 50% nucleotide identity and/or at least 65% amino acid identity and in still other instances will share at least 60% nucleotide identity and/or at least 75% amino acid identity. Watson-Crick complements of the foregoing nucleic acids also are embraced by the invention.

20 In screening for Tat-Stimulatory Factor proteins, a Southern blot may be performed using the foregoing conditions, together with a radioactive probe. After washing the membrane to which the DNA is finally transferred, the membrane can be placed against x-ray film to detect the radioactive signal.

The invention also includes degenerate nucleic acids which include alternative codons to
25 those present in the native materials. For example, serine residues are encoded by the codons TCA, AGT, TCC, TCG, TCT and AGC. Each of the six codons is equivalent for the purposes of encoding a serine residue. Thus, it will be apparent to one of ordinary skill in the art that any of the serine-encoding nucleotide triplets may be employed to direct the protein synthesis apparatus, *in vitro* or *in vivo*, to incorporate a serine residue. Similarly, nucleotide sequence triplets which
30 encode other amino acid residues include, but are not limited to,: CCA, CCC, CCG and CCT (proline codons); CGA, CGC, CGG, CGT, AGA and AGG (arginine codons); ACA, ACC, ACG and ACT (threonine codons); AAC and AAT (asparagine codons); and ATA, ATC and ATT

(isoleucine codons). Other amino acid residues may be encoded similarly by multiple nucleotide sequences. Thus, the invention embraces degenerate nucleic acids that differ from the biologically isolated nucleic acids in codon sequence due to the degeneracy of the genetic code.

The invention also provides isolated unique fragments of SEQ ID NO:1 or compliments
5 of SEQ ID NO:1. A unique fragment is one that is a 'signature' for the larger nucleic acid. It, for example, is long enough to assure that its precise sequence is not found in molecules outside of the Tat-Stimulatory Factor proteins defined above. Unique fragments can be used as probes in Southern blot assays to identify such proteins, or can be used in amplification assays such as those employing PCR. As known to those skilled in the art, large probes such as 200 BP or more
10 are preferred for certain uses such as Southern blots, while smaller fragments will be preferred for uses such as PCR. Unique fragments also can be used to produce fusion proteins for generating antibodies as demonstrated in the Examples, or for generating immunoassay components. Likewise, unique fragments can be employed to produce nonfused fragments of the Tat-Stimulatory Factor proteins, useful, for example, in immunoassays or as a competitive
15 binding partner of the kinase which binds to the Tat-Stimulatory Factor proteins, for example, in therapeutic applications. Unique fragments further can be used as antisense molecules to inhibit the expression of Tat-Stimulatory Factor proteins, particularly for therapeutic purposes as described in greater detail below.

As will be recognized by those skilled in the art, the size of the unique fragment will
20 depend upon its conservancy in the genetic code. Thus, some regions of SEQ ID NO:1 and its complement will require longer segments to be unique while others will require only short segments, typically between 12 and 32 BP (e.g. 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 bases long). Virtually any segment of the region of SEQ ID NO:1 beginning at nucleotide 53 and ending at nucleotide 2703, or its complement, that is 18 or
25 more nucleotides in length will be unique except that the unique fragments herein include consecutive nucleotides of SEQ ID NO:1 other than those nucleotides which code for SEQ ID NO:3. Those skilled in the art are well versed in methods for selecting such sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from non-Tat-Stimulatory Factor proteins. A comparison of the sequence of the
30 fragment to those on known data bases typically is all that is necessary, although *in vitro* confirmatory hybridization and sequencing analysis may be performed.

The invention also provides isolated, functional unique fragments of SEQ ID NO:2. Such sequences are useful, for example, alone or as fusion proteins to generate antibodies, as a components of an immunoassay, as an inhibitor of Tat-Stimulatory Factor activity, as a binding partner of Tat-Stimulatory Factor binding kinases (for example, for isolating such kinases) or for
5 inhibiting binding of such kinases to Tat-Stimulatory Factor proteins. Such unique fragments can be identified by routine assays, such as those involving testing a fragment's ability to generate antibodies if injected into a proper host, testing the fragment's ability to inhibit Tat-Stimulatory Factor activity, as described below, etc. A unique fragment of a Tat-Stimulatory Factor protein, in general, has the features and characteristics of unique fragments as discussed
10 above in connection with nucleic acids.

As mentioned above, the invention embraces antisense oligonucleotides that selectively bind to a nucleic acid molecule encoding a Tat-Stimulatory Factor protein, to decrease transcription activity, and in particular transcriptional elongation. This is desirable in virtually any medical condition wherein a reduction in transcriptional elongation is desirable, including to
15 reduce HIV-1 transcriptional elongation by Tat. Antisense molecules, in this manner, can be used to slow down or arrest the propagation of HIV *in vivo*.

As used herein, the term "antisense oligonucleotide" or "antisense" describes an oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological
20 conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that mRNA. The antisense molecules are designed so as to interfere with transcription or translation of a target gene upon hybridization with the target gene. Those skilled in the art will recognize that the exact length of the antisense oligonucleotide and its degree of complementarity with its target
25 will depend upon the specific target selected, including the sequence of the target and the particular bases which comprise that sequence. It is preferred that the antisense oligonucleotide be constructed and arranged so as to bind selectively with the target under physiological conditions, i.e., to hybridize substantially more to the target sequence than to any other sequence in the target cell under physiological conditions. Based upon SEQ ID NO:1, or upon allelic or
30 homologous genomic and/or cDNA sequences, one of skill in the art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. In order to be sufficiently selective and potent for inhibition, such antisense

oligonucleotides should comprise at least 10 and, more preferably, at least 15 consecutive bases which are complementary to the target. Most preferably, the antisense oligonucleotides comprise a complementary sequence of 20-30 bases. Although oligonucleotides may be chosen which are antisense to any region of the gene or mRNA transcripts, in preferred embodiments the antisense oligonucleotides correspond to N-terminal or 5' upstream sites such as translation initiation, transcription initiation or promoter sites. In addition, 3'-untranslated regions may be targeted. Targeting to mRNA splicing sites has also been used in the art but may be less preferred if alternative mRNA splicing occurs. In addition, the antisense is targeted, preferably, to sites in which mRNA secondary structure is not expected (see, e.g., Sainio et al., Cell Mol. Neurobiol. 14(5):439-457 (1994)) and at which proteins are not expected to bind. Finally, although, SEQ ID NO:1 discloses a cDNA sequence, one of ordinary skill in the art may easily derive the genomic DNA corresponding to the cDNA of SEQ ID NO:1. Thus, the present invention also provides for antisense oligonucleotides which are complementary to the genomic DNA corresponding to SEQ ID NO:1. Similarly, antisense to allelic or homologous cDNAs and genomic DNAs are enabled without undue experimentation.

In one set of embodiments, the antisense oligonucleotides of the invention may be composed of "natural" deoxyribonucleotides, ribonucleotides, or any combination thereof. That is, the 5' end of one native nucleotide and the 3' end of another native nucleotide may be covalently linked, as in natural systems, via a phosphodiester internucleoside linkage. These oligonucleotides may be prepared by art recognized methods which may be carried out manually or by an automated synthesizer. They also may be produced recombinantly by vectors.

In preferred embodiments, however, the antisense oligonucleotides of the invention also may include "modified" oligonucleotides. That is, the oligonucleotides may be modified in a number of ways which do not prevent them from hybridizing to their target but which enhance their stability or targeting or which otherwise enhance their therapeutic effectiveness.

The term "modified oligonucleotide" as used herein describes an oligonucleotide in which (1) at least two of its nucleotides are covalently linked via a synthetic internucleoside linkage (i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide) and/or (2) a chemical group not normally associated with nucleic acids has been covalently attached to the oligonucleotide. Preferred synthetic internucleoside linkages are phosphorothioates, alkylphosphonates, phosphorodithioates,

phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, and carboxymethyl esters.

The term "modified oligonucleotide" also encompasses oligonucleotides with a covalently modified base and/or sugar. For example, modified oligonucleotides include oligonucleotides having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified oligonucleotides may include a 2'-O-alkylated ribose group. In addition, modified oligonucleotides may include sugars such as arabinose instead of ribose. The present invention, thus, contemplates pharmaceutical preparations containing modified antisense molecules that are complementary to and hybridizable with, under physiological conditions, nucleic acids encoding Tat-Stimulatory Factor proteins or kinases that bind to Tat-Stimulatory Factor proteins, together with pharmaceutically acceptable carriers.

Antisense oligonucleotides may be administered as part of a pharmaceutical composition. Such a pharmaceutical composition may include the antisense oligonucleotides in combination with any standard physiologically and/or pharmaceutically acceptable carriers which are known in the art. The compositions should be sterile and contain a therapeutically effective amount of the antisense oligonucleotides in a unit of weight or volume suitable for administration to a patient. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue, or organism. The characteristics of the carrier will depend on the route of administration. Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art.

The invention also involves expression vectors coding for Tat-Stimulatory Factor proteins and fragments and variants thereof and Tat Stimulatory Factor antisense, and host cells containing those expression vectors. Virtually any cells, prokaryotic or eukaryotic, which can be transformed with heterologous DNA or RNA and which can be grown or maintained in culture, may be used in the practice of the invention. Examples include bacterial cells such as *E. coli* and mammalian cells such as mouse, hamster, pig, goat, primate, etc. They may be of a wide variety of tissue types, including mast cells, fibroblasts, oocytes and lymphocytes, and they may be primary cells or cell lines. Specific examples include CHO cells and COS cells. Cell-free

transcription systems also may be used in lieu of cells. In gene therapy applications, human hematopoietic cells that are precursors of T-cells are contemplated.

As used herein, a "vector" may be any of a number of nucleic acids into which a desired sequence may be inserted by restriction and ligation for transport between different genetic environments or for expression in a host cell. Vectors are typically composed of DNA although RNA vectors are also available. Vectors include, but are not limited to, plasmids and phagemids. A cloning vector is one which is able to replicate in a host cell, and which is further characterized by one or more endonuclease restriction sites at which the vector may be cut in a determinable fashion and into which a desired DNA sequence may be ligated such that the new recombinant vector retains its ability to replicate in the host cell. In the case of plasmids, replication of the desired sequence may occur many times as the plasmid increases in copy number within the host bacterium or just a single time per host before the host reproduces by mitosis. In the case of phage, replication may occur actively during a lytic phase or passively during a lysogenic phase. An expression vector is one into which a desired DNA sequence may be inserted by restriction and ligation such that it is operably joined to regulatory sequences and may be expressed as an RNA transcript. Vectors may further contain one or more marker sequences suitable for use in the identification of cells which have or have not been transformed or transfected with the vector. Markers include, for example, genes encoding proteins which increase or decrease either resistance or sensitivity to antibiotics or other compounds, genes which encode enzymes whose activities are detectable by standard assays known in the art (e.g. β -galactosidase or alkaline phosphatase), and genes which visibly affect the phenotype of transformed or transfected cells, colonies or plaques. Preferred vectors are those capable of autonomous replication and expression of the structural gene products present in the DNA segments to which they are operably joined.

As used herein, a coding sequence and regulatory sequences are said to be "operably" joined when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory sequences. If it is desired that the coding sequences be translated into a functional protein, two DNA sequences are said to be operably joined if induction of a promoter in the 5' regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere

with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably joined to a coding sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

5 The precise nature of the regulatory sequences needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 5' non-transcribing and 5' non-translating sequences involved with the initiation of transcription and translation respectively, such as a TATA box, capping sequence, CAAT sequence, and the like. Especially, such 5' non-transcribing regulatory sequences will include a promoter region which includes a
10 promoter sequence for transcriptional control of the operably joined gene. Regulatory sequences may also include enhancer sequences or upstream activator sequences as desired. The vectors of the invention may optionally include 5' leader or signal sequences, 5' or 3'. The choice and design of an appropriate vector is within the ability and discretion of one of ordinary skill in the art.

Expression vectors containing all the necessary elements for expression are commercially
15 available and known to those skilled in the art. See Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, 1989. Cells are genetically engineered by the introduction into the cells of heterologous DNA (RNA) encoding the Tat-Stimulatory Factor protein or fragment or variant thereof. That heterologous DNA (RNA) is placed under operable control of transcriptional elements to permit the expression of
20 the heterologous DNA in the host cell.

Examples of systems for mRNA expression in mammalian cells are those such as pRc/CMV (available from Invitrogen, Carlsbad, CA) that contain a selectable marker such as a gene that confers G418 resistance (which facilitates the selection of stably transfected cell lines) and the human cytomegalovirus (CMV) enhancer-promoter sequences. Another system suitable
25 for expression in primate or canine cell lines is the pCEP4 vector (Invitrogen), which contains an Epstein Barr virus (EBV) origin of replication, facilitating the maintenance of plasmid as a multicopy extrachromosomal element.

A variety of systems for expression of proteins in bacterial, yeast, or insect cells have been described and are commercially available. Examples of such systems include the
30 Glutathione-S-transferase (GST) Gene Fusion system available from Pharmacia Biotech, Piscataway, NJ. In this system a plasmid is constructed containing the protein sequence of interest (in this case, the transporter including the first extracellular domain) inserted in frame

downstream of the 25 kDa GST domain from S. japonicum. Expression of the fusion protein can be induced in transfected bacterial cells and the fusion protein purified by affinity chromatography using Glutathione Sepharose 4B. Cleavage of the desired peptide from the GST sequences is achieved using a site specific protease whose recognition sequence is located immediately upstream from the cloning site. An alternative system which is desirable since it maintains eukaryotic-specific functions such as glycosylation is recombination into baculovirus. Standard protocols exist (c.f. O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, IRL/Oxford University Press, 1992) and vectors, cells, and reagents are commercially available. Vaccinia virus vectors also may be employed.

10 The invention also permits the construction of Tat-Stimulatory Factor gene "knock-outs" in cells and in animals, providing materials for studying transcription and HIV replication.

 The invention also involves polypeptides which bind to Tat-Stimulatory Factor proteins, complexes of Tat-Stimulatory Factor proteins and their kinase binding partners, and to the kinase binding partners of the Tat-Stimulatory Factor proteins. Such binding partners can be used, for example, in screening assays to detect the presence or absence of Tat-Stimulatory Factor proteins and their kinase binding partners and in purification protocols to isolate Tat-Stimulatory Factor proteins and their kinase binding partners. Such poly peptides also can be used to inhibit the native activity of the Tat-Stimulatory Factor proteins or their kinase binding partners, for example, by binding to such proteins, or their binding partners or both.

20 The invention, therefore, involves antibodies or fragments of antibodies having the ability to selectively bind to Tat-Stimulatory Factor proteins. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

 Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')₂ fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody

molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also provides for F(ab')₂, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')₂ fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies.

Thus, the invention involves polypeptides of numerous size and type that specifically to Tat-Stimulatory Factor proteins, their kinase binding partners and compounds both Tat-Stimulatory Factor proteins and their kinase binding partners. These polypeptides can be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries. Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptoids and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention. Briefly, one prepares a phage library (using e.g. m13, fd, or lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional procedures. The inserts may represent, for example, a completely degenerate or biased array. One then can select phage-bearing inserts which bind to the Tat-Stimulatory Factor protein. This process can be repeated through several cycles of reselection of phage that bind to the Tat-Stimulatory Factor protein. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequence analysis can be conducted to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to the Tat-Stimulatory Factor protein can be determined. One can repeat the procedure using a biased library containing inserts containing part or all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Yeast 2 hybrid screening methods also may be used to identify polypeptides that bind to the Tat-Stimulatory Factor proteins. Thus, the Tat-Stimulatory Factor molecule of the invention, or a fragment thereof, can be used to screen peptide libraries, including phage display libraries, to identify and select peptide binding partners of the Tat-Stimulatory Factor proteins of the invention. Such molecules can be used, as described, for screening assays, for purification protocols, for interfering directly with the functioning of Tat and for other purposes that will be apparent to those of ordinary skill in the art.

The Tat-Stimulatory Factor proteins also can be used to isolate their native binding partners, including the kinases that complex with the Tat-Stimulatory Factor proteins. Such isolation of kinases may be according to well-known methods. For example, isolated Tat-Stimulatory Factor proteins can be attached to a substrate, and then a solution suspected of containing the kinase may be applied to the substrate. If the kinase binding partner for Tat-Stimulatory Factor proteins is present in the solution, then it will bind to the substrate-bound Tat-

Stimulatory Factor protein. The kinase then may be isolated. The kinase also can be isolated by successive fractionation of a solution containing the kinase, and determining whether the kinase is present with each successive phase of fractionation. Kinase activity may be determined from *in vitro* kinase reaction using Tat-Stimulatory Factor as the kinase substrate.

5 When used therapeutically, the compounds of the invention are administered in therapeutically effective amounts. In general, a therapeutically effective amount means that amount necessary to delay the onset of, inhibit the progression of, or halt altogether the particular condition being treated. Therapeutically effective amounts specifically will be those which desirably influence transcriptional activity. When it is desired to decrease such activity, then any
10 inhibition of such activity is regarded as a therapeutically effective amount. When it is desired to increase such activity, then any enhancement of such activity is regarded as a therapeutically effective amount. Generally, a therapeutically effective amount will vary with the subject's age, condition, and sex, as well as the nature and extent of the disease in the subject, all of which can be determined by one of ordinary skill in the art. The dosage may be adjusted by the individual
15 physician or veterinarian, particularly in the event of any complication. A therapeutically effective amount typically varies from 0.01 mg/kg to about 1000 mg/kg, preferably from about 0.1 mg/kg to about 200 mg/kg and most preferably from about 0.2 mg/kg to about 20 mg/kg, in one or more dose administrations daily, for one or more days.

 The therapeutics of the invention can be administered by any conventional route,
20 including injection or by gradual infusion over time. The administration may, for example, be oral, intravenous, intraperitoneal, intramuscular, intracavity, intrarespiratory, subcutaneous, or transdermal.

 Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol,
25 polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on
30 Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

The invention also contemplates gene therapy. The procedure for performing *ex vivo* gene therapy is outlined in U.S. Patent 5,399,346 and in exhibits submitted in the file history of that patent, all of which are publicly available documents. In general, it involves introduction *in vitro* of a functional copy of a gene or fragment thereof into a cell(s) of a subject and returning
5 the genetically engineered cell(s) to the subject. The functional copy of the gene or fragment thereof is under operable control of regulatory elements which permit expression of the gene in the genetically engineered cell(s). Numerous transfection and transduction techniques as well as appropriate expression vectors are well known to those of ordinary skill in the art, some of which are described in PCT application WO95/00654.

10 As an illustrative example, primary human blood cells which are precursors of T-cells can be obtained from the bone marrow of a subject who is a candidate for such gene therapy. Then, such cells can be genetically engineered *ex vivo* with DNA (RNA) encoding an agent that binds to a Tat-Stimulatory Factor nucleic acid or expression product thereof. The genetically engineered cells then are returned to the patient. Such recombinant cells are expected to resist
15 intracellular HIV replication.

The invention also contemplates targeting to particular cells the nucleic acids and proteins of the invention, including specifically antisense nucleic acids and agents that bind Tat-Stimulatory Factor proteins. Targeting may be tissue-specific, using targeting agents known to those of ordinary skill in the art. Methodologies for targeting include conjugates, such as
20 those described in U.S. Patent 5,391,723 to Priest. Another example of a well-known targeting vehicle is liposomes. Liposomes are commercially available from Gibco BRL (Life Technologies, Inc., Gaithersburg, MD). Numerous methods are published for making targeted liposomes. A preferred cell type is a T-cell, and, in particular, a T-cell infected with HIV.

According to another aspect of the invention, a method of screening for compounds
25 which bind to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase is provided.

The methods disclosed herein are useful for identifying specific compounds or molecules which bind to Tat-SF1, Tat-SF1 associated kinase or a complex of Tat-SF1 and Tat-SF1 associated kinase. Such compounds can have utility as reagents for affinity purification, for
30 example where the compound is immobilized and used to "capture" Tat-SF1, Tat-SF1 associated kinase or the complex of Tat-SF1 and Tat-SF1 associated kinase from a biological extract such as a cell or tissue homogenate. Such compounds can also be used for localization of Tat-SF1,

Tat-SF1 associated kinase or a complex of Tat-SF1 associated kinase in an intact biological system such as a cell or a tissue. The methods disclosed herein are also useful for preparation of a "library" of high probability drug candidates. Compounds can be identified as binding Tat-SF1, Tat-SF1 associated kinase or complex of Tat-SF1 associated kinase by a change in a
5 selected biological activity, such as modulation of Tat-SF1-mediated transcriptional activity, Tat-SF1 associated kinase phosphorylation activity and the like. High probability drug candidates are those compounds which cause a change in a biological activity. Such changes in activity can provide potential therapeutic benefits. Compounds identified by in vitro assays as potential drug candidates can be tested subsequently in cell- or animal-based disease models to determine more
10 accurately the therapeutic potential of the drug candidates.

One screening method is an *in vitro* binding assay of a type familiar to one of ordinary skill in the art. To preform such an assay, a test compound is contacted with the Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase and the binding of the compound to the Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1
15 and Tat-SF1-associated kinase is determined. The binding can be determined in a number of ways. For example, binding of a labeled compound can be detected by methods well known to those of ordinary skill in the art. Binding of a compound also can be determined by a competitive binding assay, such as by a reduction in binding of an antibody to the Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase or by the
20 inhibition of binding between Tat-SF1 and Tat-SF1 associated kinase. Binding of a compound also can be determined by a change in electrophoretic mobility or chromatographic elution profile of Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase relative to the profile of such a polypeptide (or complex) not bound by the compound.

Preferably, the compound is an oligonucleotide. Oligonucleotides useful in the invention
25 can be prepared according to standard methods in the art. Oligonucleotides which bind to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase are preferably prepared by binding and screening for binding activity, followed by random or targeted mutation of the nucleotides which constitute the oligonucleotide in an iterative fashion. Commercially available libraries can be screened or, as would be more likely, can be prepared
30 for screening. Preparation of peptide libraries are described herein. Lam (*Nature* 354:82-84, 1991) also describes combinatorial methods for creating libraries of synthesized peptides on polystyrene beads, each bead carrying only one peptide. Similar procedures are known for

making libraries of oligonucleotides. Methods for the selection of several classes of molecules (including oligonucleotides, peptides, and RNAs) also are described in Abelson, *Science* 249:488-489, 1990; Ellington et al., *Nature* 346:818-822, 1990; Tuerk et al., *Science* 249: 505-510, 1990; Irvine et al., *J. Mol. Biol.* 222:739-761, 1991; Bock et al., *Nature* 355:564-566, 1992; 5 Ellington et al., *Nature* 355:850-852, 1992; Gallop et al., *J. Med. Chem.* 37:1233-1251, 1994; Gordon et al., *J. Med. Chem.* 37:1385-1401, 1994; and Gold, *J. Biol. Chem.* 270:13581-13584, 1995.

In one embodiment of the assay, the compound is detectably labeled. The binding of the labeled compound bound to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and 10 Tat-SF1-associated kinase then can be determined by any method known to one of skill in the art. The particular method chosen to detect the labeled compound will depend on the nature of the label. For instance, a radioactively labeled compound can be detected by scintillation counting, autoradiography, and phosphorimaging. Fluorescently labeled compounds can be detected by fluorometry. Other detectable labels are known in the art, along with suitable 15 detection methods.

The compound also can be labeled with a molecule which serves as a binding point for a detectable label. For example, a compound can be labeled with a biotin molecule which serves as an binding point for a streptavidin molecule which is detectably labeled.

Other preferred methods for determining the binding of a compound include detecting a 20 change in a biological activity of the Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase. Determinable biological activities include Tat-SF1 mediated transcription from TAR elements, changes in protein conformation and/or protein-protein interaction which are detectable in an immunoassay, e.g. by antibody binding, Tat-SF1-associated kinase substrate phosphorylation, and binding of Tat-SF1 to a TAR element. For 25 example, Tat-SF1 mediated transcription can be determined by a reconstituted transcription assay in which Tat-SF1 is combined with transcription factors required to initiated and elongate transcription of a RNA containing a TAR element. The change in transcription mediated by Tat-SF1 which is caused by binding of a compound can be readily determined by comparing the transcription in the presence and in the absence of the compound. Tat-SF1-associated kinase 30 substrate phosphorylation can be determined by inclusion of radiolabeled ATP in a phosphorylation reaction and determined by observing the difference in the radiolabel incorporated into a substrate of the Tat-SF1-associated kinase. Modulation of binding of

Tat-SF1 to a TAR element by binding of a compound to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase can be determined by a well-known assay such as an electrophoretic mobility shift assay (EMSA). A change in electrophoretic mobility observed upon contacting the compound with Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase will reflect a change in the binding to a TAR element. Changes in protein conformation and/or protein-protein interaction are detectable by immunoassays using antibodies which recognize a particular protein conformation or protein-protein interaction such as the interaction of Tat-SF1 and Tat-SF1-associated kinase. A change in antibody binding subsequent to contacting Tat-SF1, Tat-SF1-associated kinase and/or a complex of the two with a compound is readily determined using standard immunoassay techniques. Immunoassays which measure disruption of antibody binding to a particular epitope of Tat-SF1, Tat-SF1-associated kinase and/or a complex of the two are useful for determining binding of a compound to that particular epitope. One of ordinary skill in the art will recognize that a panel of antibodies which recognize a variety of epitopes will enable determination of the binding site of a compound which binds to Tat-SF1, Tat-SF1-associated kinase and/or a complex of the two. Biological activities and methods of determining a change in the activities subsequent to binding of a compound to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase, are described more fully in the Examples below.

It is not intended that the foregoing represents an exhaustive listing of methods useful for determining the binding of a compound to Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase. Additional detectable labels, or biological activities of Tat-SF1, Tat-SF1-associated kinase, or a complex of Tat-SF1 and Tat-SF1-associated kinase will be apparent to one of ordinary skill in the art.

According to another aspect of the invention, a method of screening for compounds which modulate Tat-SF1-mediated transcriptional activity is provided. The method involves (1) providing a mammalian cell containing a gene encoding a Tat-SF1 polypeptide; (2) contacting the mammalian cell with one or more compounds, preferably under conditions to induce Tat-SF1 mediated transcription; and (3) determining the Tat-SF1 mediated transcriptional activation.

Cell-based screening methods are provided herein for determining the Tat-SF1-mediated transcriptional activity modulating potential of compounds. These methods are useful for identifying compounds which can modulate Tat-SF1-mediated transcriptional activity in the presence of cellular proteins and other factors involved in the transcription of genes in a cell.

Thus, such methods permit screening of compounds in a variety of cells which may be particularly relevant to a disease state. For example, to identify compounds which are useful for reducing Tat-SF1-mediated transcriptional activity as a means of reducing HIV-1 infection, one of ordinary skill in the art can select an appropriate cell within the host range of HIV-1, such as a T cell, to perform the screening assay in.

The skilled artisan can readily determine the effect of a test compound on Tat-SF1-mediated transcriptional activation in a cell-based assay. It is known that the presence of Tat-SF1 in a cell stimulates transcription of genes which are operably linked to a TAR sequence, in particular by increasing elongation of nascent transcripts. Thus detection of a change in Tat-SF1 mediated-transcriptional activation involves detecting increased (or decreased) transcription of a TAR-linked nucleic acid. The skilled artisan can choose a nucleic acid sequence to link to a TAR element, operably link the TAR element to the nucleic acid using standard molecular biology techniques, and test for a change in transcription in nucleic acid by standard techniques, e.g., quantitative nucleic acid hybridization, polymerase chain reaction amplification, nuclease protection assay and the like. Alternatively, one of ordinary skill in the art can use a nucleic acid which contains a TAR element in the foregoing assay to determine the effect of a test compound on Tat-SF1-mediated transcriptional activity. For example, HIV-1 sequences containing a TAR element can be used in such an assay, such as a reporter construct containing HIV-1 LTR linked to the bacterial CAT gene. Preferably an indicator gene is transcribed and the amount of indicator gene product is determined in the presence and the absence of the compound.

In preferred embodiments, the mammalian cell used in the assay of Tat-SF1-mediated transcriptional activation modulating activity also contains a gene encoding a Tat polypeptide and/or an indicator gene operably linked to a TAR element and/or a gene encoding a Tat-SF1-associated kinase polypeptide. These genes can be included in a variety of mammalian expression vectors, including plasmid- or virus-based episomal vectors and vectors which integrate into the host cell chromosomes, as is known to those skilled in the art. The genes described above can be introduced into the mammalian cell by standard procedures, or can be resident in the cell, such as encoded by the cell's chromosomal nucleic acid. Standard procedures for introducing nucleic acids into a mammalian cell include, but are not limited to, physical means such as transfection, electroporation and bombardment with nucleic acid-coated microparticles, and biological means such as receptor-mediated endocytosis using targeted nucleic acids or nucleic acids contained in targeted liposomes and viral infection. The particular

vectors and/or means of introducing genes may depend on the mammalian cell chosen for the assay. The cells and cell lines useful in such assays include HeLa cells, COS cells and the like.

Of course, any of the foregoing genes can include only part of the gene and/or encode only a part of the polypeptide where the gene product to be detected is a nucleic acid, or where
5 the portion of the polypeptide encoded by the gene retains the activity of the whole polypeptide or an epitope bound by an antibody. For example, an indicator gene need only encode a RNA gene product that is detectable by hybridization in an assay such as RNase protection, oligonucleotide hybridization, or reverse transcriptase polymerase chain reaction (RT-PCR). Where a polypeptide gene product is to be detected, a fragment which is sufficient to retain a
10 detectable activity in an immunoassay, or a enzymatic activity assay, would be sufficient to meet the criteria disclosed above.

As used herein, modulation of Tat-SF1-mediated transcriptional activation refers to the ability of a compound to modulate Tat transcriptional activation, mediated by Tat-SF1, from a TAR RNA element. Thus, a Tat-SF1-mediated transcriptional activation modulating activity
15 refers to the ability of a compound to reduce or increase the ability of Tat-SF1 to stimulate Tat-dependent transcriptional elongation of RNA molecules containing a TAR element. For example, compounds which have Tat-SF1-mediated transcriptional activation modulating activity, include compounds which disrupt the binding of Tat-SF1 to a TAR element, compounds which disrupt the binding of Tat-SF1 to a Tat-SF1-associated kinase, compounds which disrupt
20 the phosphorylation of Tat-SF1 by a Tat-SF1-associated kinase, compound which disrupt the recruitment of additional transcription factors to the RNA containing a TAR element, and the like.

Tat and Tat-SF1 proteins can bind to a TAR element in a promoter which controls the transcription of an indicator gene. The indicator gene product, whether nucleic acid or protein,
25 provides a readily detectable output for screening compounds for Tat-SF1-mediated transcriptional activation modulating activity. An indicator gene permits high throughput screening of a number of compounds at one time. Depending on the choice of indicator gene, and its gene product, automation of the methods disclosed herein is also contemplated.

Preferred indicator genes encode an indicator gene product which is easily detectable
30 with or without disruption of the mammalian cell. Gene products such as RNA transcripts of the gene can be detected by hybridization assays or any other assay which detects nucleic acids of a specific sequence. Preferably, hybridization assays are conducted under stringent conditions as

defined above. Indicator gene products which are proteins can be detected by any technique suitable for detection of proteins. For example, an indicator gene can encode an indicator gene product to which antibodies have been raised. Such antibodies which selectively bind to the indicator gene product can be employed in immunoassays to determine the amount of protein
5 produced as a result of increased Tat-responsive transcription in the presence and absence of a compound. Antibodies useful in the detection of indicator gene products include commercially available antibodies such as antibodies to green fluorescent protein (Clontech, Palo Alto, CA), *E. coli* bacterial alkaline phosphatase, β -galactosidase (Boehringer Mannheim, Indianapolis, IN), and luciferase (Cortex Biochemicals, San Leandro, CA).

10 The indicator gene product can be a protein which has an enzymatic activity that can be assayed. Methods for determining the amount of such enzymes by colorimetric means (for example, conversion of X-gal into a blue product by β -galactosidase) or radioactive means (for example, addition of a ^3H -acetyl or ^{14}C -acetyl group to a chloramphenicol molecule by chloramphenicol acetyl transferase) will be known to one of ordinary skill in the art. Other
15 indicator gene products, such as green fluorescent protein, are directly detectable by colorimetric means.

Preferably, the indicator gene is selected from the group consisting of the genes encoding the following proteins: β -galactosidase, alkaline phosphatase, chloramphenicol acetyl transferase, luciferase, and green fluorescent protein. However, the skilled artisan may select any
20 indicator gene for use in accordance with the methods of the invention provided that the indicator gene product is detectable.

The indicator gene is operably linked to a promoter containing at least one TAR element which drives expression of the indicator gene by serving as the locus for binding of Tat, Tat-SF1, associated proteins and a RNA polymerase.

25 The nucleic acid molecules introduced into the mammalian cell may be contained on a plasmid or other extrachromosomal nucleic acid, or may be incorporated into the cell's chromosomes. Non-chromosomal nucleic acids include plasmids, phagemids, bacteriophage genomes, virus genomes, and the like. Non-chromosomal nucleic acids useful for preparation of expression vectors are well known in the art and thus are not described further here.

30

Examples

Example 1: Tat activation of HIV-1 transcription requires a specific cellular activity. Tat-SF

We have previously established a partially reconstituted transcription reaction that supports a Tat-specific and TAR-dependent activation of HIV transcription (Zhou and Sharp, *EMBO J.* 14: 321-328, 1995; Figure 1A). This reaction requires a Tat-SF (Tat-Stimulatory Factor) activity that is specific for Tat stimulation of elongation. It also requires a phosphocellulose 0.5-1.0 M KOAc fraction of HeLa nuclear extract, termed the pc-D fraction, and the purified basal factors TFIID, TFIIA, and transcription factor Sp1. Reactions containing these components, but lacking Tat-SF activity supported activation by Sp1 and GAL4-VP16 (Zhou and Sharp, 1995), but not by Tat (lanes 1 and 2, Figure 1A). The transcription reactions were performed as follows. Reconstituted transcription reactions containing both templates pHIV+TAR-G400 and pHIVTAR-G100 were performed in the absence (-) or presence (+) of the 0.5 M KCl Q-Sepharose chromatographic fraction containing Tat-SF activity as described previously (Zhou and Sharp, 1995). The pc-D fraction and purified TFIIA, TFIID, and Sp1 were present in all reactions. G-less cassettes of two different lengths were inserted into the above two templates at position +955 downstream of the HIV-1 initiation site to measure the effect of Tat on transcriptional elongation. Transcripts derived from these two templates were digested by RNase T1 and the resulting 400- and 100-nucleotide G-less RNA fragments were separated in a denaturing polyacrylamide gel, and their positions were indicated by the arrows. In the presence of a partially purified Tat-SF fraction, Tat specifically increased the number of transcripts elongating beyond 1000 nucleotides from a HIV-1 promoter containing the wild-type TAR element (pHIV+TAR-G400), but not from an internal control promoter with a mutant TAR (pHIVTAR-G100, compare lanes 3 and 4).

The pc-D fraction was shown by Western blotting to contain the basal transcription factors TFIIB, IIE, IIF, IIH, and RNA polymerase II (Zhou and Sharp, 1995). This fraction probably also contains other activities important for Tat function, because it can not be substituted with highly purified basal transcription factors. Using the reconstituted reaction detailed above to follow the activity, Tat-SF was further purified by several chromatographic steps. HeLa nuclear extract in buffer D/0.1 M KCl (20 mM Hepes-KOH, pH 7.9/20% (vol/vol) glycerol/0.1 M KCl/0.2 mM EDTA/0.5 mM dithiothreitol/0.5 mM PMSF) was loaded on a phosphocellulose column preequilibrated with the same buffer. The flowthrough was loaded on a DEAE-Sepharose FF (Pharmacia, Piscataway, NJ) matrix column preequilibrated with buffer D/0.1 M KCl. After washing the column with the same buffer, Tat-SF activity was eluted from the column with buffer D/0.3 M KCl. This fraction was dialyzed against buffer D/0.1 M KCl

and applied to a Q-Sepharose FF (Pharmacia) matrix column preequilibrated with the same buffer. The column was washed with buffer D/0.1 M KCl and the bound proteins were eluted with a 0.1-0.7 M KCl gradient made in buffer D. Fractions were analyzed for Tat-SF activity in reconstituted transcription assays and for pp140 in kinase reactions (as described below). The
5 0.4-0.5 M KCl Q-Sepharose fraction containing Tat-SF activity and pp140 was dialyzed against buffer D/0.1 M KCl and applied to a Heparin Sepharose column. After washing the column extensively with buffer D/0.1 M KCl, Tat-SF/pp140 was eluted with increasing salt concentrations and was found mostly in 0.2-0.4 M KCl fractions. These fractions were combined, dialyzed to 0.1 M KCl, and loaded on a Glutathione Sepharose (Pharmacia) column
10 containing GST-Tat fusion proteins. After washing with buffer D/0.4 M KCl, Tat-SF/pp140 was eluted from the column with buffer D containing 1.4 M KCl. The estimated overall purification after these steps was ~3000-fold.

Example 2: Detection of a complex containing Tat, a cellular kinase and a 140 kD
15 phosphoprotein in the transcription reaction.

Phosphorylation of RNA polymerase II has been implicated in regulation of the processivity of elongation (O'Brien et al., *Nature* 370:75-77, 1994; Dahmus, *Biochim Biophys Acta* 1261:171-182, 1995). To investigate whether protein phosphorylation might be associated with Tat-SF, an immobilized HIV TAR RNA was used to collect bound proteins from a
20 reconstituted transcription reaction in the presence of γ -³²P ATP. Referring to Figure 1B, biotinylated TAR RNA (nucleotides +1 to +82) immobilized on the Paramagnetic beads was introduced into the kinase reactions containing γ -³²P ATP (10 mCi) and either the pc-D fraction alone (lanes 3 and 4), or the Tat-SF fraction alone (lanes 5 and 6), or both pc-D and Tat-SF fractions together (lanes 1, 2, and 7). Recombinant wild type Tat protein (13 ng) was included in
25 the reactions shown in lanes 2, 4, and 6. Tat mutant TatK41A (13 ng) was present in lane 7 (Rice and Carlotti, *J. Virol* 64:1864-1868, 1990). After incubation for 10 min at 30°C, the TAR RNA beads were washed extensively in buffer D containing 100 mM KCl and 0.1% NP-40 and the bound proteins were analyzed by SDS-PAGE. In reactions containing either the pc-D fraction alone (lanes 3 and 4, Figure 1B) or the Tat-SF fraction alone (lanes 5 and 6), addition of
30 Tat did not consistently affect the phosphorylation of proteins on immobilized TAR. When both fractions were incubated together in the presence of Tat, phosphorylation of a protein of approximately 140 kD, termed pp140, was observed (Figure 1B, lane 2). In the absence of Tat,

however, only a small amount of phosphorylated pp140 was detected (lanes 1). Thus, collection of a phosphorylated pp140 on TAR required the presence of the pc-D fraction, the Tat-SF fraction, and Tat. This suggested the existence of a complex on TAR that consists of Tat, a cellular kinase probably derived from the pc-D fraction, and the kinase substrate pp140 from the
5 Tat-SF fraction (see below).

An intact Tat activation domain was necessary for pp140 phosphorylation on TAR. When a non-functional Tat mutant (K41A; Rice and Carlotti, 1990), which has the lysine at position 41 substituted by alanine, was present in the kinase reaction (Figure 1B, lane 7), the amount of phosphorylated pp140 collected on the immobilized TAR was significantly reduced (compare
10 lanes 2 and 7). Importantly, this was not due to a decreased ability of K41A to interact with TAR, since K41A bound to TAR as efficiently as wild type Tat in a gel mobility-shift assay.

Similar results were also obtained with another Tat mutant (TatC), which lacks the cysteine-rich activation domain (amino acids 22 to 37; Rice and Carlotti, 1990) and is completely defective for transcriptional activity (Figure 1C). Kinase reactions containing an
15 immobilized TAR were prepared as above. Recombinant wild type Tat protein (13 ng) or Tat mutant (TatC, 13 ng) lacking the cysteine-rich domain (amino acid 22-37) was included in the reactions as indicated. No Tat was present in the control reaction (lane 1).

The polypeptide pp140 co-purified and co-titrated with Tat-SF transcriptional activity during purification through multiple chromatography steps. For example, when partially purified
20 Tat-SF was sedimented through a glycerol gradient and the first 10 fractions were analyzed in the reconstituted transcription assay for the presence of Tat-SF activity (Figure 2A), Tat-SF activity and pp140 co-peaked. A 0.2-0.4 M KCl Heparin Sepharose fraction (load, lanes 3 and 4), prepared as described above, containing Tat-SF activity was loaded onto a 12-35% glycerol gradient and subjected to ultracentrifugation. The first 10 of a total 16 fractions were tested for
25 Tat-SF activity in the reconstituted transcription reactions in the absence (-) or presence (+) of Tat as in Figure 1A. Control reactions (lanes 1 and 2) did not contain Tat-SF fraction. Protein molecular weight markers were sedimented in a parallel gradient and analyzed by silver staining. The peak of Tat-SF activity that supported Tat *trans*-activation was in fractions #4 and #3, corresponding to a native molecular mass of approximately 100 kD. When the same fractions
30 were analyzed in the kinase assay for the presence of pp140 in a kinase assay as described in Figure 1B (see Figure 2B), phosphorylation of pp140 was also most evident in fractions #4 and #3, in agreement with the transcription results.

Example 3: Purification of pp140 by Tat affinity column chromatography.

As noted above, detection of the phosphorylated pp140 on TAR requires the presence of the pc-D fraction, the Tat-SF fraction, and Tat. Sequential incubations of these components with an immobilized TAR revealed a stable and direct interaction between Tat and a cellular kinase and between Tat and pp140. Therefore, columns containing immobilized Tat were used to affinity-purify Tat-SF/pp140. The 0.2-0.4 M KCl Heparin Sepharose fraction (load) containing Tat-SF activity described in Example 1 was subjected to fractionation through an Affi-Gel 10 matrix column (Bio-Rad, Hercules, CA) containing immobilized Tat. Tat-SF activity was eluted from the column with increasing salt concentrations. The 0.6 M KCl fraction was analyzed in Figure 3. Fractions eluted from either a GST-Tat column or a Tat affinity column were analyzed in reconstituted transcription assays for the presence of Tat-SF activity (Figure 3A). Both fractions were enriched in Tat-SF activity which supported a Tat-specific and TAR-dependent activation. The same two fractions were tested in a kinase assay as described in Example 1 and were found to contain pp140 (Figure 3B). When analyzed by silver staining, the polypeptide profiles of these two fractions were different overall, with the only common band having a mobility of 140 kD (Figure 3C). This polypeptide was judged to be pp140 and probably a component of Tat-SF activity.

The 140 kD polypeptide was recovered from the SDS polyacrylamide gel by blotting onto a nitrocellulose membrane. Approximately 15 μ g pp140 was recovered from the membrane and subjected to digestion with lys-C. Six major peptides were obtained and microsequenced. Sequence analysis of six peptides indicated that pp140 was a novel protein. However, one of the peptides (KMNAQETATGMAFEETIDE, SEQ ID NO:3) was contained in the sequence of an unidentified "expressed sequence tag" (EST) EST60354 in the Washington University/Merck EST database. A 103 amino acid protein fragment (Figure 5A, amino acids 387 to 489) encoded by the corresponding EST clone was expressed as a GST fusion and used to immunize rabbits for the production of polyclonal antisera. By Western blotting, the affinity-purified antibody specifically recognized a 140 kD protein present in both HeLa nuclear extracts and a partially purified Tat-SF fraction (Figure 4C).

Example 4: pp140 and a cellular kinase form a complex independently of Tat.

The Tat-SF specific antibody was used to test the relationship between the EST clone and pp140. The Tat-SF fraction was subjected to immunodepletion with the affinity purified antibody and then incubated together with the pc-D fraction and Tat. The polypeptide pp140 was immunodepleted from a 0.4-0.5 M KCl Q-Sepharose fraction (Tat-SF fraction) through
5 incubation on ice twice for 1.5 hr each with the affinity purified anti-pp140 antibody immobilized on protein A Sepharose beads. Referring to Fig. 4, the depleted (lanes 5 and 6) or undepleted Tat-SF fractions (no depl. lanes 1-4) were incubated with the pc-D fraction in the absence (-) or presence (+) of Tat and the reactions were subjected to immunoprecipitation with the immobilized anti-pp140 antibody (lanes 3-6). Preimmune antibody was used in control
10 precipitations (lanes 1 and 2). An unfractionated HeLa nuclear extract (NE) with (+) or without (-) the addition of Tat was also subjected to immunoprecipitation with the specific antibody (lanes 7 and 8). After extensive washes with buffer D containing 100 mM KCl, 0.1% NP-40, and 10 mM MgCl₂, the immune-complex bound to protein A Sepharose beads was analyzed in a kinase reaction in the presence of γ -³²P ATP for 10 min at 30°C, washed with buffer D, and
15 analyzed by SDS-PAGE (Figure 4A, lane 6). In contrast to the control undepleted Tat-SF fraction (lane 4), the fraction depleted with the specific antibody did not contain the phosphorylated pp140 (compare lanes 4 and 6). Therefore, the 140 kD protein recovered from the SDS gel and represented by the EST clone was indeed pp140, the kinase substrate.

These reactions also suggest that the polypeptide pp140 and its kinase formed a stable
20 complex independently of Tat. When the Tat-SF fraction and the pc-D fraction were incubated together in the absence of Tat, followed by immunoprecipitation with the anti-pp140 antibody, pp140 was phosphorylated by its associated kinase when the isolated immune-complex was assayed in the kinase reaction (Figure 4A, lane 3). This result indicates that pp140 forms a complex with its kinase in the absence of Tat. Furthermore, the addition of Tat to the initial
25 incubation did not change the level of phosphorylation on pp140 (compare lanes 3 and 4).

A preformed complex containing pp140 and its kinase could be isolated by immunoprecipitation and detected in a kinase reaction from an unfractionated HeLa nuclear extract in the absence of Tat (Figure 4A, compare lanes 7 and 8). This complex was stable under transcription conditions (less than 0.1 M KCl), but dissociated in washes of greater than 0.25 M
30 KCl, and probably dissociated during fractionation in the purification of Tat-SF. These observations suggest that Tat is not required for the phosphorylation of pp140 by its associated kinase, but is required for the association of the phosphorylated pp140 and the kinase with TAR

(Figure 1B). Thus, Tat probably recruits a preformed complex containing pp140 and a kinase to the HIV promoter region during transcription.

Example 5: pp140 is required for Tat-SF transcriptional activity.

5 To test whether pp140 is indeed necessary for Tat activation, the anti-pp140 antibody was used to immuno-deplete pp140 from a partially purified fraction containing Tat-SF activity and the depleted fraction was then tested in reconstituted transcription reactions for its ability to support Tat activation (Figure 4B). A 0.4-0.5 M KCl Q-Sepharose fraction containing Tat-SF activity was subjected to immunodepletion with preimmune antibody (lanes 5, 6, 9, and 10) or
10 specific anti-pp140 antibody (lanes 7, 8, 11, and 12) immobilized on protein A Sepharose beads as in Example 4. Tat-SF fraction subjected to depletion once (1X) or twice (2X), and the undepleted Tat-SF fraction (lanes 3 and 4) were tested in transcription reactions for Tat-SF activity as described above. No Tat-SF fraction was present in the control reactions (lanes 1 and 2). Control reactions without Tat-SF did not support Tat activation (Figure 4B, lanes 1 and 2).
15 Inclusion of the Tat-SF fraction resulted in a TAR-dependent activation by Tat as expected (lanes 3 and 4). As compared to control Tat-SF fraction (lanes 3 and 4), depletion of the Tat-SF fraction with specific anti-pp140 antibody immobilized on protein A Sepharose matrix either once (lanes 7 and 8) or especially twice (lanes 11 and 12) significantly reduced its ability to support Tat activation. In contrast, depletion of the Tat-SF fraction with preimmune antibody
20 either once (lanes 5 and 6) or twice (lanes 9 and 10) did not significantly reduce Tat activation. Similarly, depletion of the Tat-SF fraction with an unrelated antibody (anti-HA mAb 12CA5) had no effect Tat activation.

The undepleted Tat-SF fraction and Tat-SF fraction twice-depleted with the specific or preimmune antibody were subjected to electrophoresis and Western blotting with the anti-pp140
25 antisera. As expected, a Western blot (Figure 4C) indicates that depletion with the anti-pp140 antibody efficiently removed pp140 from Tat-SF fraction. Taken together, these experiments strongly argue that pp140 is indeed necessary for Tat-SF transcriptional activity.

Example 6: Isolation of the cDNA encoding pp140.

30 An XhoI-EcoRI fragment ³²P-labeled DNA probe made from the insert of the EST clone corresponding to the COOH-terminus of the Tat-SF1 gene and its 3' untranslated region was labeled and used as a probe to screen a λZipLox (Gibco BRL) cDNA library prepared from

human HL60 cells (provided by J. Borrow, MIT, Cambridge, MA). cDNAs were recovered from seven independent plaques in the autonomously-replicating plasmid pZL1 using the protocol provided by the manufacturer (Gibco BRL). Inserts from the seven independent plaques had similar restriction endonuclease cleavage patterns, and sequencing confirmed that they contained overlapping segments. The largest cDNA clone containing the full length Tat-SF1 gene was named pZL-Tat-SF1-4b and was sequenced by dideoxy-DNA sequencing with T7 DNA polymerase. The largest cDNA fragment was 2.8-kb in length and contained a 2271-bp open reading frame. There were multiple in-frame stop codons both upstream and downstream of this coding region. Surprisingly, the open reading frame encoded a protein of 754 amino acids with a calculated molecular weight of 85,767 daltons (Figure 5A), which was significantly less than the apparent molecular weight of 140 kD calculated from the mobility in an SDS polyacrylamide gel.

This cDNA was judged to encode the authentic full length pp140 based on several observations. First, transfection of this cDNA into human 293T cells (Pear et al., *Proc. Natl. Acad. Sci. U S A* 90:8392-8396, 1993) resulted in the production of the full length pp140 polypeptide and thus a significant increase in the total cellular level of pp140 as judged by Western blotting. Second, all six peptide sequences obtained from partial sequencing of pp140 were found in the predicted coding region (underlined in Figure 5A). Third, Northern analysis of poly(A)⁺ RNA isolated from several different types of human cells detected a single 3.0 kb species, a length consistent with that of the cDNA segment and adequate to encode a polypeptide of 86 kD. Finally, this cDNA and two additional cDNA clones isolated from a completely different cDNA library had identical upstream in-frame stop codons.

Sequence analysis of the protein, referred to as Tat-SF1, is shown in Figure 5A.

Glutamate (E) and aspartate (D) residues present in the COOH-terminal half of Tat-SF1 (amino acids 420 to 754) are shown in bold face type. The two RNA recognition motifs (RRMs) in the NH₂-terminal half of Tat-SF1 are boxed, with the conserved RNP1 and RNP2 motifs shown in shaded area and bold face type, respectively. The six peptides of Tat-SF1 that were generated by digestion with lys-C and subjected to microsequencing are underlined. The regions of Tat-SF1 that are homologous to human EWS are underlined with broken lines.

The sequence analysis of the protein revealed that it has several unique features. The protein can be roughly divided at position 420 into two halves. The COOH-terminal half was extremely rich in acidic amino acids, with 48% of the last 245 amino acid residues as glutamate

or aspartate. The unusual acidic nature of this protein may be responsible for its aberrant mobility in an SDS gel. The COOH-terminal half also contained many serine residues that are arranged in a short peptide sequence matching consensus sites for phosphorylation by Casein Kinase II (Marshak and Carroll, *Methods Enzymol.* 200:134-156, 1991). Such phosphorylation would contribute more negative charges to this region.

The NH₂-terminal half of Tat-SF1 contained two tandem RNA recognition motifs (Kenan et al., *Trends Biochem. Sci.* 16:214-220, 1991) which have homology to many RNA-binding proteins. Interestingly, the first RRM of Tat-SF1 (amino acid 128 to 217, boxed in Figure 5A) was similar in length and displayed the strongest sequence homology to the RRMs located in the COOH-terminal half of two closely related human proteins, EWS (Delattre et al., *Nature* 359:162-165, 1992; Sorensen et al., *Nature Genet.* 6:146-151, 1994) (Figure 5B) and FUS/TLS (Croizat et al., *Nature* 363:640-644, 1993; Rabbitts et al., *Nature Genet.* 4:175-180, 1993). Furthermore, the sequence homology between Tat-SF1 and EWS, or between Tat-SF1 and FUS/TLS, extended beyond the two RRMs into the immediate NH₂-terminal region of Tat-SF1 (Figure 5A and B). The amino acid sequences of the homologous regions of Tat-SF1 (SEQ ID NO:2) and EWS (SEQ ID NO:4) are compared in Fig. 5B. The amino acids of each protein are numbered next to the sequences. Vertical lines and dots indicate identical and conserved residues, respectively. EWS has two tandem, imperfect repeats (amino acids 209 to 236) that show homology to Tat-SF1 (amino acids 30 to 44). The alignment between the first repeat (amino acids 209-223) of EWS and Tat-SF1 is shown. The first RRM of Tat-SF1 (amino acids 128 to 217) is almost identical in length, and is 27% identical and 52% similar in amino acid sequence to the RRM of EWS. Sequence homology similar to that observed between Tat-SF1 and EWS also exists between Tat-SF1 and human FUS/TLS, which is closely related to EWS. The RRMs of other RNA binding proteins are less homologous and show greater variations in length as revealed by the BLAST algorithm (Altschul et al., *J. Mol. Biol.* 215:403-410, 1990).

These observations suggest that Tat-SF1 is related to EWS and FUS/TLS, which are members of a novel class of putative transcription factors that presumably interact with RNA. Both EWS and FUS/TLS are involved in many forms of human solid tumors (Ladanyi, *Diagn. Mol. Pathol.* 4:162-173, 1995; Rabbitts, *Nature* 372:143-149, 1994), such as Ewing's sarcoma (Delattre et al., 1992; Sorensen et al., 1994) and human myxoid liposarcoma (Croizat et al., 1993; Rabbitts et al., 1993), through chromosomal translocations.

Example 7: Overexpression of Tat-SF1 enhances Tat activation in vivo.

To investigate whether overexpression of Tat-SF1 affects the level of Tat activation *in vivo*, a plasmid expressing Tat-SF1 and a reporter construct containing HIV-1 LTR linked to the bacterial CAT gene were introduced into HeLa cells either in the presence or absence of a
5 cotransfected plasmid expressing Tat (Table I and Figure 6). As a control, the effect of Tat-SF1 overexpression on transcriptional activation by the acidic activation domain VP16 in the TFEB-VP16 and GAL4-VP16 fusion proteins was assayed (Harper et al., *Proc. Natl. Acad. Sci. U S A* 93:8536-8540, 1996). The Tat-SF1 gene was subcloned into the mammalian expressing vector pSV7d (Truett et al., *DNA* 4:333-349, 1985) to create pSV-Tat-SF1. pSV-Tat-SF1 or vector
10 pSV7d and a reporter construct pBennCAT (Gendelman et al., *Proc. Natl. Acad. Sci. U S A* 83:9759-63, 1986) containing HIV-1 LTR linked to the bacterial CAT gene (1 μ g each) and an internal control plasmid pCMV β -Gal were co-transfected into HeLa cells, either in the presence or absence of a Tat expressing plasmid pcTat (0.3 μ g) (Tiley et al., *Virology* 178:560-567, 1990). CAT activity was measured 48 hr later as described (Neumann et al., *BioTechniques* 5:444-447,
15 1987). In control experiments, pSV-Tat-SF1 or pSV7d and the reporter construct pMyc3E1BLuc (Harper et al., 1996) were introduced into HeLa cells together with the plasmids pRCCMV-TFEB-VP16 (0.3 μ g) expressing the TFEB-VP16 fusion protein (Harper et al., 1996). pMyc3E1BLuc contained the luciferase gene downstream of the Adenovirus E1B promoter with three binding sites for TFEB. Reporter construct pG5E1BCAT (Lillie et al., *Nature* 338:39-44,
20 1989) containing five GAL4-binding sites inserted upstream of the E1B promoter and the CAT gene was used to assay GAL4-VP16 *trans*-activation. The fold activation by Tat or VP16 in cells containing the empty vector was assigned a value of 1, and activation in the presence of Tat-SF1 was adjusted accordingly. The mean value from three experiments was shown.

Expression of Tat-SF1 from the transfected DNA consistently resulted in an increase in
25 Tat activation by an average of 5.2-fold as compared to the control HeLa cells transfected with an empty vector (Figure 6 and Table I). The enhanced activation mediated by Tat-SF1 was Tat-specific, since overexpression of Tat-SF1 had little, or sometimes even a slightly negative, effect on transcriptional activation by TFEB-VP16. Interestingly, the elevated fold induction by Tat resulting from Tat-SF1 overexpression was caused by a combination of a decrease in the basal
30 level of transcription from HIV-1 LTR in the absence of Tat and a small increase in the level of Tat-activated transcription (Table I). Since Tat-SF1 is probably a component of a protein complex that also includes a cellular kinase and perhaps other cellular components,

overexpression of Tat-SF1 alone may disrupt the normal stoichiometry of the complex resulting in a decrease in the basal level of HIV transcription. The presence of Tat could stabilize and recruit the active form of the complex to the HIV promoter to stimulate the processivity of elongation.

5

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

10

All references disclosed herein are incorporated by reference in their entirety.

A Sequence Listing is presented below and is followed by what is claimed.

SEQUENCE LISTING

(1) GENERAL INFORMATION

5

(i) APPLICANT:

- (A) NAME: MASSACHUSETTS INSTITUTE OF TECHNOLOGY
- (B) STREET: 77 MASSACHUSETTS AVENUE
- (C) CITY: CAMBRIDGE
- (D) STATE: MASSACHUSETTS
- (E) COUNTRY: UNITED STATES OF AMERICA
- (F) POSTAL CODE: 02139

10

(i) APPLICANT/INVENTOR:

- (A) NAME: SHARP, PHILLIP A.
- (B) STREET: 36 FERMONT AVENUE
- (C) CITY: NEWTON
- (D) STATE: MASSACHUSETTS
- (E) COUNTRY: UNITED STATES OF AMERICA
- (F) POSTAL CODE: 02158

15

20

(i) APPLICANT/INVENTOR:

- (A) NAME: ZHOU, QIANG
- (B) STREET: 206 STANLEY HALL
- (C) CITY: BERKELEY
- (D) STATE: CALIFORNIA
- (E) COUNTRY: UNITED STATES OF AMERICA
- (F) POSTAL CODE: 94720

25

30

- (ii) TITLE OF THE INVENTION: TAT-SF: COFACTOR FOR STIMULATION
OF TRANSCRIPTIONAL ELONGATION BY HIV-1 TAT

- (iii) NUMBER OF SEQUENCES: 5

35

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: Wolf, Greenfield & Sacks, P.C.
- (B) STREET: 600 Atlantic Avenue
- (C) CITY: Boston
- (D) STATE: MA
- (E) COUNTRY: U.S.A.
- (F) ZIP: 02210

40

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Diskette
- (B) COMPUTER: IBM Compatible
- (C) OPERATING SYSTEM: DOS
- (D) SOFTWARE: FastSEQ for Windows

45

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:

5

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 60/021,218
- (B) FILING DATE: 03-JUL-1996

10

- (A) APPLICATION NUMBER: US 60/033,152
- (B) FILING DATE: 13-DEC-1996

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Gates, Edward R.
- (B) REGISTRATION NUMBER: 31,616
- (C) REFERENCE/DOCKET NUMBER: M0656/7024WO

15

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 617-720-3500
- (B) TELEFAX: 617-720-2441

20

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2815 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

30

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 110...2371
- (D) OTHER INFORMATION:

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGGAAAGCTG GTACGCCTGC AGGTACCGGT CCGGAATTCC GGCCGNGTCG AAAGCGTCAT 60
40 TTGGGCTCT TAGTTCTTCT GAACCTGCT OCTGAGCTAG GTAGGAAAC ATG AGC GGC 118
Met Ser Gly
1

ACC AAC TTG GAT GGG AAC GAT GAG TTT GAT GAG CAG TTG CGA ATG CAA 166
45 Thr Asn Leu Asp Gly Asn Asp Glu Phe Asp Glu Gln Leu Arg Met Gln
5 10 15

	GAA TTG TAC GGA GAC GGC AAG GAT GGT GAC ACC CAG ACC GAT GCC GGC	214
	Glu Leu Tyr Gly Asp Gly Lys Asp Gly Asp Thr Gln Thr Asp Ala Gly	
	20 25 30 35	
5	GGA GAA CCC GAT TCT CTC GGG CAG CAG CCG ACG GAC ACT CCC TAC GAG	262
	Gly Glu Pro Asp Ser Leu Gly Gln Gln Pro Thr Asp Thr Pro Tyr Glu	
	40 45 50	
10	TGG GAC CTG GAC AAA AAG GCT TGG TTC CCC AAG ATT ACT GAA GAT TTC	310
	Trp Asp Leu Asp Lys Lys Ala Trp Phe Pro Lys Ile Thr Glu Asp Phe	
	55 60 65	
15	ATT GCT ACA TAT CAG GCC AAT TAT GGC TTC TCT AAC GAT GGC GCA TCT	358
	Ile Ala Thr Tyr Gln Ala Asn Tyr Gly Phe Ser Asn Asp Gly Ala Ser	
	70 75 80	
20	AGT TCT ACC GCA AAT GTT GAA GAT GTC CAT GCT AGG ACT GCA GAG GAA	406
	Ser Ser Thr Ala Asn Val Glu Asp Val His Ala Arg Thr Ala Glu Glu	
	85 90 95	
25	CCT CCA CAA GAA AAA GCC CCG GAA CCC ACT GAT GCC AGA AAG AAG GGA	454
	Pro Pro Gln Glu Lys Ala Pro Glu Pro Thr Asp Ala Arg Lys Lys Gly	
	100 105 110 115	
30	GAA AAA AGA AAG GCT GAG TCA GGA TGG TTT CAT GTT GAA GAA GAC AGA	502
	Glu Lys Arg Lys Ala Glu Ser Gly Trp Phe His Val Glu Glu Asp Arg	
	120 125 130	
35	AAT ACA AAT GTA TAC GTG TCT GGT TTG CCT CCA GAT ATT ACA GTG GAT	550
	Asn Thr Asn Val Tyr Val Ser Gly Leu Pro Pro Asp Ile Thr Val Asp	
	135 140 145	
40	GAA TTT ATA CAA CTT ATG TCC AAG TTT GGC ATT ATT ATG AGA GAT CCT	598
	Glu Phe Ile Gln Leu Met Ser Lys Phe Gly Ile Ile Met Arg Asp Pro	
	150 155 160	
45	CAG ACA GAA GAA TTT AAG GTC AAA CTT TAC AAA GAT AAT CAA GGA AAT	646
	Gln Thr Glu Glu Phe Lys Val Lys Leu Tyr Lys Asp Asn Gln Gly Asn	
	165 170 175	
50	CTT AAA GGA GAC GGT CTT TGC TGT TAT TTG AAA AGA GAA TCT GTG GAA	694
	Leu Lys Gly Asp Gly Leu Cys Cys Tyr Leu Lys Arg Glu Ser Val Glu	
	180 185 190 195	
55	CTT GCA TTA AAA CTT TTG GAT GAA GAT GAA ATT AGA GGC TAC AAA TTA	742
	Leu Ala Leu Lys Leu Leu Asp Glu Asp Glu Ile Arg Gly Tyr Lys Leu	
	200 205 210	

	CAT GTT GAG GTG GCA AAG TTT CAA CTG AAG GGA GAA TAT GAT GCC TCA	790
	His Val Glu Val Ala Lys Phe Gln Leu Lys Gly Glu Tyr Asp Ala Ser	
	215 220 225	
5	AAG AAG AAG AAG AAG TGC AAA GAC TAT AAG AAG AAG CTG TCT ATG CAA	838
	Lys Lys Lys Lys Lys Cys Lys Asp Tyr Lys Lys Lys Leu Ser Met Gln	
	230 235 240	
10	CAA AAG CAG TTG GAT TGG AGA CCT GAG AGG CGA GCC GGA CCA TCC OGG	886
	Gln Lys Gln Leu Asp Trp Arg Pro Glu Arg Arg Ala Gly Pro Ser Arg	
	245 250 255	
15	ATG CGC CAT GAG CGA GTT GTC ATC ATC AAG AAT ATG TTT CAT CCT ATG	934
	Met Arg His Glu Arg Val Val Ile Ile Lys Asn Met Phe His Pro Met	
	260 265 270 275	
20	GAT TTT GAG GAT GAT CCG TTG GTG CTG AAT GAG ATC AGA GAA GAC CTT	982
	Asp Phe Glu Asp Asp Pro Leu Val Leu Asn Glu Ile Arg Glu Asp Leu	
	280 285 290	
25	CGA GTA GAG TGT TCG AAG TTT GGA CAA ATT AGG AAA CTC CTT CTC TTT	1030
	Arg Val Glu Cys Ser Lys Phe Gly Gln Ile Arg Lys Leu Leu Leu Phe	
	295 300 305	
30	GAT AGG CAC CCA GAT GGT GTG GCC TCT GTG TCC TTT CCG GAT CCA GAG	1078
	Asp Arg His Pro Asp Gly Val Ala Ser Val Ser Phe Arg Asp Pro Glu	
	310 315 320	
35	GAA GCT GAT TAT TGT ATT CAG ACT CTC GAT GGA AGA TGG TTT GGT GGC	1126
	Glu Ala Asp Tyr Cys Ile Gln Thr Leu Asp Gly Arg Trp Phe Gly Gly	
	325 330 335	
40	CGT CAA ATC ACT GCC CAG GCA TGG GAT GGG ACT ACA GAT TAT CAG GTG	1174
	Arg Gln Ile Thr Ala Gln Ala Trp Asp Gly Thr Thr Asp Tyr Gln Val	
	340 345 350 355	
45	GAG GAA ACC TCA AGA GAA AGG GAG GAA AGG CTG AGA GGA TGG GAG GCT	1222
	Glu Glu Thr Ser Arg Glu Arg Glu Glu Arg Leu Arg Gly Trp Glu Ala	
	360 365 370	
50	TTC CTC AAT GCT CCT GAG GCC AAC AGA GGC CTT AGC GTT CAG ATT CTG	1270
	Phe Leu Asn Ala Pro Glu Ala Asn Arg Gly Leu Ser Val Gln Ile Leu	
	375 380 385	
55	TCT CTG CTT CGA AAG GCA GGG CCT TCT AGA GCA AGG CAT TTT TCA GAG	1318
	Ser Leu Leu Arg Lys Ala Gly Pro Ser Arg Ala Arg His Phe Ser Glu	
	390 395 400	

	CAC CCC AGC ACA TCT AAA ATG AAT GCT CAA GAA ACT GCA ACT GGA ATG	1366
	His Pro Ser Thr Ser Lys Met Asn Ala Gln Glu Thr Ala Thr Gly Met	
	405 410 415	
5	GCA TTT GAA GAA CCT ATA GAT GAG AAG AAG TTT GAA AAG ACA GAA GAT	1414
	Ala Phe Glu Glu Pro Ile Asp Glu Lys Lys Phe Glu Lys Thr Glu Asp	
	420 425 430 435	
10	GGG GGA GAA TTT GAA GAA GGT GCT TCT GAA AAC AAT GCT AAG GAA AGT	1462
	Gly Gly Glu Phe Glu Glu Gly Ala Ser Glu Asn Asn Ala Lys Glu Ser	
	440 445 450	
15	AGC CCC GAA AAA GAG GCT GAA GAA GGC TGC CCT GAA AAA GAA TCT GAA	1510
	Ser Pro Glu Lys Glu Ala Glu Glu Gly Cys Pro Glu Lys Glu Ser Glu	
	455 460 465	
20	GAG GGC TGC CCC AAA AGA GGG TTT GAA GGC AGC TGC TCC CAA AAA GAG	1558
	Glu Gly Cys Pro Lys Arg Gly Phe Glu Gly Ser Cys Ser Gln Lys Glu	
	470 475 480	
25	TCT GAA GAA GGC AAT CCC GTA AGA GGA TCT GAA GAG GAT AGT CCT AAA	1606
	Ser Glu Glu Gly Asn Pro Val Arg Gly Ser Glu Glu Asp Ser Pro Lys	
	485 490 495	
30	AAA GAG TCT AAA AAG AAG ACA CTC AAA AAT GAT TGT GAA GAG AAT GGC	1654
	Lys Glu Ser Lys Lys Lys Thr Leu Lys Asn Asp Cys Glu Glu Asn Gly	
	500 505 510 515	
35	CTT GCA AAG GAA TCT GAA GAT GAC CTC AAC AAG GAG TCT GAA GAG GAG	1702
	Leu Ala Lys Glu Ser Glu Asp Asp Leu Asn Lys Glu Ser Glu Glu Glu	
	520 525 530	
40	GTT GGC CCC ACA AAA GAG TCC GAA GAA GAT GAC TCA GAG AAA GAG TCT	1750
	Val Gly Pro Thr Lys Glu Ser Glu Glu Asp Asp Ser Glu Lys Glu Ser	
	535 540 545	
45	GAT GAA GAC TGC TCT GAA AAA CAG TCT GAA GAT GGC TCC GAA AGA GAA	1798
	Asp Glu Asp Cys Ser Glu Lys Gln Ser Glu Asp Gly Ser Glu Arg Glu	
	550 555 560	
50	TTT GAA GAA AAT GGT CTC GAG AAA GAT TTG GAC GAG GAA GGT TCT GAA	1846
	Phe Glu Glu Asn Gly Leu Glu Lys Asp Leu Asp Glu Glu Gly Ser Glu	
	565 570 575	
55	AAG GAG CTT CAT GAA AAT GTT CTT GAC AAA GAG TTA GAA GAA AAT GAC	1894
	Lys Glu Leu His Glu Asn Val Leu Asp Lys Glu Leu Glu Glu Asn Asp	
	580 585 590 595	

	TCT GAA AAC TCC GAA TTT GAA GAT GAC GGC TCT GAA AAA GTG TTA GAT	1942
	Ser Glu Asn Ser Glu Phe Glu Asp Asp Gly Ser Glu Lys Val Leu Asp	
	600 605 610	
5	GAG GAA GGC TCT GAG AGA GAG TTT GAC GAA GAT TCA GAT GAA AAG GAA	1990
	Glu Glu Gly Ser Glu Arg Glu Phe Asp Glu Asp Ser Asp Glu Lys Glu	
	615 620 625	
10	GAA GAG GAG GAT ACA TAT GAA AAA GTA TTT GAT GAT GAG TCT GAT GAG	2038
	Glu Glu Glu Asp Thr Tyr Glu Lys Val Phe Asp Asp Glu Ser Asp Glu	
	630 635 640	
15	AAA GAG GAT GAA GAA TAT GCA GAT GAA AAG GGG CTT GAA GCT GCT GAT	2086
	Lys Glu Asp Glu Glu Tyr Ala Asp Glu Lys Gly Leu Glu Ala Ala Asp	
	645 650 655	
20	AAA AAG GCG GAA GAA GGT GAT GCA GAT GAA AAG CTG TTT GAA GAG TCA	2134
	Lys Lys Ala Glu Glu Gly Asp Ala Asp Glu Lys Leu Phe Glu Glu Ser	
	660 665 670 675	
	GAT GAC AAG GAA GAT GAA GAT GCA GAT GGA AAG GAA GTT GAA GAT GCT	2182
	Asp Asp Lys Glu Asp Glu Asp Ala Asp Gly Lys Glu Val Glu Asp Ala	
	680 685 690	
25	GAC GAA AAG TTG TTC GAA GAT GAT GAT TCC AAT GAG AAG TTG TTT GAT	2230
	Asp Glu Lys Leu Phe Glu Asp Asp Asp Ser Asn Glu Lys Leu Phe Asp	
	695 700 705	
30	GAG GAG GAA GAT TCC AGT GAG AAG TTG TTT GAC GAT TCT GAT GAG AGG	2278
	Glu Glu Glu Asp Ser Ser Glu Lys Leu Phe Asp Asp Ser Asp Glu Arg	
	710 715 720	
35	GGG ACT TTG GGT GGT TTT GGG AGT GTT GAA GAA GGG CCC CTA TCC ACT	2326
	Gly Thr Leu Gly Gly Phe Gly Ser Val Glu Glu Gly Pro Leu Ser Thr	
	725 730 735	
40	GGC AGC AGC TTT ATT CTC AGT AGC GAT GAT GAT GAC GAT GAT ATT TAATC	2376
	Gly Ser Ser Phe Ile Leu Ser Ser Asp Asp Asp Asp Asp Ile	
	740 745 750	
45	CCTTAAACTT GCTTTT TAGG GAGAGTCCTC CATCTACATT TGCCTGTGCT TCAGGGTAAT	2436
	TACTAGTAGT GTTACATGAA CATGTGCATA GTGGTAGGAT GCCATCAGAT TAAAGCAATTG	2496
	AAGTGTTC TTTGTTACCTG TACCTAATGG TTTTAAATAT ATGTTAATTG ATTGTTTAGT	2556
	TAAAATGTCA TAGTTACAAT GCAAGTAAAC TGGATACCTG TTCTTTTGTC AGATTGTGTA	2616
	AATGCATGCA GAATAATATT TTTAAGAGTA TTGATTGAAG TTTGTGATAT TCATCAATAA	2676
	AAATGAGTTG ATAATATGCA GAAACTGAAA AAAAAAAAAA AAAAAAAGT CGACNCGGCC	2736
	GGAATCCCG GGTGACGAG CTCCTAGTC GCGCGCGCT CTAGAGGATC CAAGCTTACG	2796

TACGCGTGCA TGGGACGTC

2815

(2) INFORMATION FOR SEQ ID NO:2:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 754 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

15 Met Ser Gly Thr Asn Leu Asp Gly Asn Asp Glu Phe Asp Glu Gln Leu
   1           5           10           15
Arg Met Gln Glu Leu Tyr Gly Asp Gly Lys Asp Gly Asp Thr Gln Thr
   20           25           30
Asp Ala Gly Gly Glu Pro Asp Ser Leu Gly Gln Gln Pro Thr Asp Thr
20   35           40           45
Pro Tyr Glu Trp Asp Leu Asp Lys Lys Ala Trp Phe Pro Lys Ile Thr
   50           55           60
Glu Asp Phe Ile Ala Thr Tyr Gln Ala Asn Tyr Gly Phe Ser Asn Asp
65           70           75           80
25 Gly Ala Ser Ser Ser Thr Ala Asn Val Glu Asp Val His Ala Arg Thr
   85           90           95
Ala Glu Glu Pro Pro Gln Glu Lys Ala Pro Glu Pro Thr Asp Ala Arg
   100          105          110
Lys Lys Gly Glu Lys Arg Lys Ala Glu Ser Gly Trp Phe His Val Glu
30   115          120          125
Glu Asp Arg Asn Thr Asn Val Tyr Val Ser Gly Leu Pro Pro Asp Ile
   130          135          140
Thr Val Asp Glu Phe Ile Gln Leu Met Ser Lys Phe Gly Ile Ile Met
145          150          155          160
35 Arg Asp Pro Gln Thr Glu Glu Phe Lys Val Lys Leu Tyr Lys Asp Asn
   165          170          175
Gln Gly Asn Leu Lys Gly Asp Gly Leu Cys Cys Tyr Leu Lys Arg Glu
   180          185          190
Ser Val Glu Leu Ala Leu Lys Leu Leu Asp Glu Asp Glu Ile Arg Gly
40   195          200          205
Tyr Lys Leu His Val Glu Val Ala Lys Phe Gln Leu Lys Gly Glu Tyr
   210          215          220
Asp Ala Ser Lys Lys Lys Lys Lys Cys Lys Asp Tyr Lys Lys Lys Leu
225          230          235          240
45 Ser Met Gln Gln Lys Gln Leu Asp Trp Arg Pro Glu Arg Arg Ala Gly
   245          250          255
Pro Ser Arg Met Arg His Glu Arg Val Val Ile Ile Lys Asn Met Phe

```

				260					265					270			
	His	Pro	Met	Asp	Phe	Glu	Asp	Asp	Pro	Leu	Val	Leu	Asn	Glu	Ile	Arg	
			275						280					285			
	Glu	Asp	Leu	Arg	Val	Glu	Cys	Ser	Lys	Phe	Gly	Gln	Ile	Arg	Lys	Leu	
5		290					295					300					
	Leu	Leu	Phe	Asp	Arg	His	Pro	Asp	Gly	Val	Ala	Ser	Val	Ser	Phe	Arg	
	305				310						315					320	
	Asp	Pro	Glu	Glu	Ala	Asp	Tyr	Cys	Ile	Gln	Thr	Leu	Asp	Gly	Arg	Trp	
				325						330					335		
10	Phe	Gly	Gly	Arg	Gln	Ile	Thr	Ala	Gln	Ala	Trp	Asp	Gly	Thr	Thr	Asp	
				340					345					350			
	Tyr	Gln	Val	Glu	Glu	Thr	Ser	Arg	Glu	Arg	Glu	Glu	Arg	Leu	Arg	Gly	
			355					360					365				
	Trp	Glu	Ala	Phe	Leu	Asn	Ala	Pro	Glu	Ala	Asn	Arg	Gly	Leu	Ser	Val	
15		370					375					380					
	Gln	Ile	Leu	Ser	Leu	Leu	Arg	Lys	Ala	Gly	Pro	Ser	Arg	Ala	Arg	His	
	385					390					395					400	
	Phe	Ser	Glu	His	Pro	Ser	Thr	Ser	Lys	Met	Asn	Ala	Gln	Glu	Thr	Ala	
				405						410					415		
20	Thr	Gly	Met	Ala	Phe	Glu	Glu	Pro	Ile	Asp	Glu	Lys	Lys	Phe	Glu	Lys	
			420						425					430			
	Thr	Glu	Asp	Gly	Gly	Glu	Phe	Glu	Glu	Gly	Ala	Ser	Glu	Asn	Asn	Ala	
		435				440					445						
	Lys	Glu	Ser	Ser	Pro	Glu	Lys	Glu	Ala	Glu	Glu	Gly	Cys	Pro	Glu	Lys	
25		450					455					460					
	Glu	Ser	Glu	Glu	Gly	Cys	Pro	Lys	Arg	Gly	Phe	Glu	Gly	Ser	Cys	Ser	
	465					470					475					480	
	Gln	Lys	Glu	Ser	Glu	Glu	Gly	Asn	Pro	Val	Arg	Gly	Ser	Glu	Glu	Asp	
				485						490					495		
30	Ser	Pro	Lys	Lys	Glu	Ser	Lys	Lys	Lys	Thr	Leu	Lys	Asn	Asp	Cys	Glu	
			500						505					510			
	Glu	Asn	Gly	Leu	Ala	Lys	Glu	Ser	Glu	Asp	Asp	Leu	Asn	Lys	Glu	Ser	
		515						520					525				
	Glu	Glu	Glu	Val	Gly	Pro	Thr	Lys	Glu	Ser	Glu	Glu	Asp	Asp	Ser	Glu	
35		530					535					540					
	Lys	Glu	Ser	Asp	Glu	Asp	Cys	Ser	Glu	Lys	Gln	Ser	Glu	Asp	Gly	Ser	
	545					550					555					560	
	Glu	Arg	Glu	Phe	Glu	Glu	Asn	Gly	Leu	Glu	Lys	Asp	Leu	Asp	Glu	Glu	
				565					570					575			
40	Gly	Ser	Glu	Lys	Glu	Leu	His	Glu	Asn	Val	Leu	Asp	Lys	Glu	Leu	Glu	
				580					585					590			
	Glu	Asn	Asp	Ser	Glu	Asn	Ser	Glu	Phe	Glu	Asp	Asp	Gly	Ser	Glu	Lys	
		595						600					605				
	Val	Leu	Asp	Glu	Glu	Gly	Ser	Glu	Arg	Glu	Phe	Asp	Glu	Asp	Ser	Asp	
45		610					615					620					
	Glu	Lys	Glu	Glu	Glu	Glu	Asp	Thr	Tyr	Glu	Lys	Val	Phe	Asp	Asp	Glu	
	625					630					635					640	

Ser Asp Glu Lys Glu Asp Glu Glu Tyr Ala Asp Glu Lys Gly Leu Glu
645 650 655
Ala Ala Asp Lys Lys Ala Glu Glu Gly Asp Ala Asp Glu Lys Leu Phe
660 665 670
5 Glu Glu Ser Asp Asp Lys Glu Asp Glu Asp Ala Asp Gly Lys Glu Val
675 680 685
Glu Asp Ala Asp Glu Lys Leu Phe Glu Asp Asp Asp Ser Asn Glu Lys
690 695 700
Leu Phe Asp Glu Glu Glu Asp Ser Ser Glu Lys Leu Phe Asp Asp Ser
10 705 710 715 720
Asp Glu Arg Gly Thr Leu Gly Gly Phe Gly Ser Val Glu Glu Gly Pro
725 730 735
Leu Ser Thr Gly Ser Ser Phe Ile Leu Ser Ser Asp Asp Asp Asp Asp
740 745 750
15 Asp Ile

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Lys Met Asn Ala Gln Glu Thr Ala Thr Gly Met Ala Phe Glu Glu Pro
1 5 10 15
30 Ile Asp Glu

(2) INFORMATION FOR SEQ ID NO:4:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 656 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Ala Ser Thr Asp Tyr Ser Thr Tyr Ser Gln Ala Ala Ala Gln Gln
45 1 5 10 15
Gly Tyr Ser Ala Tyr Thr Ala Gln Pro Thr Gln Gly Tyr Ala Gln Thr
20 25 30

Thr Gln Ala Tyr Gly Gln Gln Ser Tyr Gly Thr Tyr Gly Gln Pro Thr
 35 40 45
 Asp Val Ser Tyr Thr Gln Ala Gln Thr Thr Ala Thr Tyr Gly Gln Thr
 50 55 60
 5 Ala Tyr Ala Thr Ser Tyr Gly Gln Pro Pro Thr Gly Tyr Thr Thr Pro
 65 70 75 80
 Thr Ala Pro Gln Ala Tyr Ser Gln Pro Val Gln Gly Tyr Gly Thr Gly
 85 90 95
 Ala Tyr Asp Thr Thr Thr Ala Thr Val Thr Thr Thr Gln Ala Ser Tyr
 10 100 105 110
 Ala Ala Gln Ser Ala Tyr Gly Thr Gln Pro Ala Tyr Pro Ala Tyr Gly
 115 120 125
 Gln Gln Pro Ala Ala Thr Ala Pro Thr Arg Pro Gln Asp Gly Asn Lys
 130 135 140
 15 Pro Thr Glu Thr Ser Gln Pro Gln Ser Ser Thr Gly Gly Tyr Asn Gln
 145 150 155 160
 Pro Ser Leu Gly Tyr Gly Gln Ser Asn Tyr Ser Tyr Pro Gln Val Pro
 165 170 175
 Gly Ser Tyr Pro Met Gln Pro Val Thr Ala Pro Pro Ser Tyr Pro Pro
 20 180 185 190
 Thr Ser Tyr Ser Ser Thr Gln Pro Thr Ser Tyr Asp Gln Ser Ser Tyr
 195 200 205
 Ser Gln Gln Asn Thr Tyr Gly Gln Pro Ser Ser Tyr Gly Gln Gln Ser
 210 215 220
 25 Ser Tyr Gly Gln Gln Ser Ser Tyr Gly Gln Gln Pro Pro Thr Ser Tyr
 225 230 235 240
 Pro Pro Gln Thr Gly Ser Tyr Ser Gln Ala Pro Ser Gln Tyr Ser Gln
 245 250 255
 Gln Ser Ser Ser Tyr Gly Gln Gln Ser Ser Phe Arg Gln Asp His Pro
 30 260 265 270
 Ser Ser Met Gly Val Tyr Gly Gln Glu Ser Gly Gly Phe Ser Gly Pro
 275 280 285
 Gly Glu Asn Arg Ser Met Ser Gly Pro Asp Asn Arg Gly Arg Gly Arg
 290 295 300
 35 Gly Gly Phe Asp Arg Gly Gly Met Ser Arg Gly Gly Arg Gly Gly Gly
 305 310 315 320
 Arg Gly Gly Met Gly Ser Ala Gly Glu Arg Gly Gly Phe Asn Lys Pro
 325 330 335
 Gly Gly Pro Met Asp Glu Gly Pro Asp Leu Asp Leu Gly Pro Pro Val
 40 340 345 350
 Asp Pro Asp Glu Asp Ser Asp Asn Ser Ala Ile Tyr Val Gln Gly Leu
 355 360 365
 Asn Asp Ser Val Thr Leu Asp Asp Leu Ala Asp Phe Phe Lys Gln Cys
 370 375 380
 45 Gly Val Val Lys Met Asn Lys Arg Thr Gly Gln Pro Met Ile His Ile
 385 390 395 400
 Tyr Leu Asp Lys Glu Thr Gly Lys Pro Lys Gly Asp Ala Thr Val Ser

				405					410					415
	Tyr	Glu	Asp	Pro	Pro	Thr	Ala	Lys	Ala	Ala	Val	Glu	Trp	Phe
				420					425					430
	Lys	Asp	Phe	Gln	Gly	Ser	Lys	Leu	Lys	Val	Ser	Leu	Ala	Arg
5			435				440						445	
	Pro	Pro	Met	Asn	Ser	Met	Arg	Gly	Gly	Leu	Pro	Pro	Arg	Glu
			450			455					460			
	Gly	Met	Pro	Pro	Pro	Leu	Arg	Gly	Gly	Pro	Gly	Gly	Pro	Gly
			465			470					475			480
10	Gly	Gly	Pro	Met	Gly	Arg	Met	Gly	Gly	Arg	Gly	Gly	Asp	Arg
					485					490				495
	Phe	Pro	Pro	Arg	Gly	Pro	Arg	Gly	Ser	Arg	Gly	Asn	Pro	Ser
				500				505					510	
	Gly	Asn	Val	Gln	His	Arg	Ala	Gly	Asp	Trp	Gln	Cys	Pro	Asn
15			515					520					525	
	Cys	Gly	Asn	Gln	Asn	Phe	Ala	Trp	Arg	Thr	Glu	Cys	Asn	Gln
			530				535						540	
	Ala	Pro	Lys	Pro	Glu	Gly	Phe	Leu	Pro	Pro	Pro	Phe	Pro	Pro
			545			550					555			560
20	Gly	Asp	Arg	Gly	Arg	Gly	Gly	Pro	Gly	Gly	Met	Arg	Gly	Gly
					565					570				575
	Gly	Leu	Met	Asp	Arg	Gly	Gly	Pro	Gly	Gly	Met	Phe	Arg	Gly
				580				585					590	
	Gly	Gly	Asp	Arg	Gly	Gly	Phe	Arg	Gly	Gly	Arg	Gly	Met	Asp
25			595					600					605	
	Gly	Phe	Gly	Gly	Gly	Arg	Arg	Gly	Gly	Pro	Gly	Gly	Pro	Pro
			610				615					620		
	Leu	Met	Glu	Gln	Met	Gly	Gly	Arg	Arg	Gly	Gly	Arg	Gly	Gly
			625			630				635				640
30	Lys	Met	Asp	Lys	Gly	Glu	His	Arg	Gln	Glu	Arg	Arg	Asp	Arg
				645						650				655

(2) INFORMATION FOR SEQ ID NO:5:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2672 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

40

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: Coding Sequence

45

(B) LOCATION: 58...2319

(D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

	AGCGTCATTT	CGGCTCTTA	GTTCTTCTGA	ACCGTGCTCC	TGAGCTAGGT	AGGAAAC	ATG	60
							Met	
5							1	
	AGC GGC ACC AAC TTG GAT GGG AAC GAT GAG TTT GAT GAG CAG TTG CGA							108
	Ser Gly Thr Asn Leu Asp Gly Asn Asp Glu Phe Asp Glu Gln Leu Arg							
		5		10		15		
10	ATG CAA GAA TTG TAC GGA GAC GGC AAG GAT GGT GAC ACC CAG ACC GAT							156
	Met Gln Glu Leu Tyr Gly Asp Gly Lys Asp Gly Asp Thr Gln Thr Asp							
		20		25		30		
15	GCC GGC GGA GAA CCC GAT TCT CTC GGG CAG CAG CCG ACG GAC ACT CCC							204
	Ala Gly Gly Glu Pro Asp Ser Leu Gly Gln Gln Pro Thr Asp Thr Pro							
		35		40		45		
	TAC GAG TGG GAC CTG GAC AAA AAG GCT TGG TTC CCC AAG ATT ACT GAA							252
20	Tyr Glu Trp Asp Leu Asp Lys Lys Ala Trp Phe Pro Lys Ile Thr Glu							
		50		55		60		65
	GAT TTC ATT GCT ACA TAT CAG GCC AAT TAT GGC TTC TCT AAC GAT GGC							300
25	Asp Phe Ile Ala Thr Tyr Gln Ala Asn Tyr Gly Phe Ser Asn Asp Gly							
			70		75		80	
	GCA TCT AGT TCT ACC GCA AAT GTT GAA GAT GTC CAT GCT AGG ACT GCA							348
	Ala Ser Ser Ser Thr Ala Asn Val Glu Asp Val His Ala Arg Thr Ala							
		85		90		95		
30	GAG GAA CCT CCA CAA GAA AAA GCC CCG GAA CCC ACT GAT GCC AGA AAG							396
	Glu Glu Pro Pro Gln Glu Lys Ala Pro Glu Pro Thr Asp Ala Arg Lys							
		100		105		110		
35	AAG GGA GAA AAA AGA AAG GCT GAG TCA GGA TGG TTT CAT GTT GAA GAA							444
	Lys Gly Glu Lys Arg Lys Ala Glu Ser Gly Trp Phe His Val Glu Glu							
		115		120		125		
	GAC AGA AAT ACA AAT GTA TAC GTG TCT GGT TTG CCT CCA GAT ATT ACA							492
40	Asp Arg Asn Thr Asn Val Tyr Val Ser Gly Leu Pro Pro Asp Ile Thr							
		130		135		140		145
	GTG GAT GAA TTT ATA CAA CTT ATG TCC AAG TTT GGC ATT ATT ATG AGA							540
45	Val Asp Glu Phe Ile Gln Leu Met Ser Lys Phe Gly Ile Ile Met Arg							
		150		155		160		
	GAT CCT CAG ACA GAA GAA TTT AAG GTC AAA CTT TAC AAA GAT AAT CAA							588

	Asp	Pro	Gln	Thr	Glu	Glu	Phe	Lys	Val	Lys	Leu	Tyr	Lys	Asp	Asn	Gln	
				165					170					175			
5	GGA	AAT	CTT	AAA	GGA	GAC	GGT	CTT	TGC	TGT	TAT	TTG	AAA	AGA	GAA	TCT	636
	Gly	Asn	Leu	Lys	Gly	Asp	Gly	Leu	Cys	Cys	Tyr	Leu	Lys	Arg	Glu	Ser	
			180					185					190				
10	GTG	GAA	CTT	GCA	TTA	AAA	CTT	TTG	GAT	GAA	GAT	GAA	ATT	AGA	GGC	TAC	684
	Val	Glu	Leu	Ala	Leu	Lys	Leu	Leu	Asp	Glu	Asp	Glu	Ile	Arg	Gly	Tyr	
		195					200					205					
15	AAA	TTA	CAT	GTT	GAG	GTG	GCA	AAG	TTT	CAA	CTG	AAG	GGA	GAA	TAT	GAT	732
	Lys	Leu	His	Val	Glu	Val	Ala	Lys	Phe	Gln	Leu	Lys	Gly	Glu	Tyr	Asp	
	210					215				220						225	
20	GCC	TCA	AAG	AAG	AAG	AAG	AAG	TGC	AAA	GAC	TAT	AAG	AAG	AAG	CTG	TCT	780
	Ala	Ser	Lys	Lys	Lys	Lys	Lys	Cys	Lys	Asp	Tyr	Lys	Lys	Lys	Leu	Ser	
				230						235					240		
25	ATG	CAA	CAA	AAG	CAG	TTG	GAT	TGG	AGA	CCT	GAG	AGG	CGA	GCC	GGA	CCA	828
	Met	Gln	Gln	Lys	Gln	Leu	Asp	Trp	Arg	Pro	Glu	Arg	Arg	Ala	Gly	Pro	
				245				250						255			
30	TCC	CGG	ATG	CGC	CAT	GAG	CGA	GTT	GTC	ATC	ATC	AAG	AAT	ATG	TTT	CAT	876
	Ser	Arg	Met	Arg	His	Glu	Arg	Val	Val	Ile	Ile	Lys	Asn	Met	Phe	His	
			260					265					270				
35	CCT	ATG	GAT	TTT	GAG	GAT	GAT	CCG	TTG	GTG	CTG	AAT	GAG	ATC	AGA	GAA	924
	Pro	Met	Asp	Phe	Glu	Asp	Asp	Pro	Leu	Val	Leu	Asn	Glu	Ile	Arg	Glu	
		275					280					285					
40	GAC	CTT	CGA	GTA	GAG	TGT	TCG	AAG	TTT	GGA	CAA	ATT	AGG	AAA	CTC	CTT	972
	Asp	Leu	Arg	Val	Glu	Cys	Ser	Lys	Phe	Gly	Gln	Ile	Arg	Lys	Leu	Leu	
	290					295				300					305		
45	CTC	TTT	GAT	AGG	CAC	CCA	GAT	GGT	GTG	GCC	TCT	GTG	TCC	TTT	CGG	GAT	1020
	Leu	Phe	Asp	Arg	His	Pro	Asp	Gly	Val	Ala	Ser	Val	Ser	Phe	Arg	Asp	
				310				315						320			
50	CCA	GAG	GAA	GCT	GAT	TAT	TGT	ATT	CAG	ACT	CTC	GAT	GGA	AGA	TGG	TTT	1068
	Pro	Glu	Glu	Ala	Asp	Tyr	Cys	Ile	Gln	Thr	Leu	Asp	Gly	Arg	Trp	Phe	
				325				330					335				
55	GGT	GGC	CGT	CAA	ATC	ACT	GCC	CAG	GCA	TGG	GAT	GGG	ACT	ACA	GAT	TAT	1116
	Gly	Gly	Arg	Gln	Ile	Thr	Ala	Gln	Ala	Trp	Asp	Gly	Thr	Thr	Asp	Tyr	
			340					345				350					

	CAG GTG GAG GAA ACC TCA AGA GAA AGG GAG GAA AGG CTG AGA GGA TGG	1164
	Gln Val Glu Glu Thr Ser Arg Glu Arg Glu Glu Arg Leu Arg Gly Trp	
	355 360 365	
5	GAG GCT TTC CTC AAT GCT CCT GAG GCC AAC AGA GGC CTT AGC GTT CAG	1212
	Glu Ala Phe Leu Asn Ala Pro Glu Ala Asn Arg Gly Leu Ser Val Gln	
	370 375 380 385	
10	ATT CTG TCT CTG CTT CGA AAG GCA GGG CCT TCT AGA GCA AGG CAT TTT	1260
	Ile Leu Ser Leu Leu Arg Lys Ala Gly Pro Ser Arg Ala Arg His Phe	
	390 395 400	
15	TCA GAG CAC CCC AGC ACA TCT AAA ATG AAT GCT CAA GAA ACT GCA ACT	1308
	Ser Glu His Pro Ser Thr Ser Lys Met Asn Ala Gln Glu Thr Ala Thr	
	405 410 415	
20	GGA ATG GCA TTT GAA GAA CCT ATA GAT GAG AAG AAG TTT GAA AAG ACA	1356
	Gly Met Ala Phe Glu Glu Pro Ile Asp Glu Lys Lys Phe Glu Lys Thr	
	420 425 430	
	GAA GAT GGG GGA GAA TTT GAA GAA GGT GCT TCT GAA AAC AAT GCT AAG	1404
	Glu Asp Gly Gly Glu Phe Glu Glu Gly Ala Ser Glu Asn Asn Ala Lys	
	435 440 445	
25	GAA AGT AGC CCC GAA AAA GAG GCT GAA GAA GGC TGC CCT GAA AAA GAA	1452
	Glu Ser Ser Pro Glu Lys Glu Ala Glu Glu Gly Cys Pro Glu Lys Glu	
	450 455 460 465	
30	TCT GAA GAG GGC TGC CCC AAA AGA GGG TTT GAA GGC AGC TGC TCC CAA	1500
	Ser Glu Glu Gly Cys Pro Lys Arg Gly Phe Glu Gly Ser Cys Ser Gln	
	470 475 480	
35	AAA GAG TCT GAA GAA GGC AAT CCC GTA AGA GGA TCT GAA GAG GAT AGT	1548
	Lys Glu Ser Glu Glu Gly Asn Pro Val Arg Gly Ser Glu Glu Asp Ser	
	485 490 495	
40	CCT AAA AAA GAG TCT AAA AAG AAG ACA CTC AAA AAT GAT TGT GAA GAG	1596
	Pro Lys Lys Glu Ser Lys Lys Lys Thr Leu Lys Asn Asp Cys Glu Glu	
	500 505 510	
	AAT GGC CTT GCA AAG GAA TCT GAA GAT GAC CTC AAC AAG GAG TCT GAA	1644
	Asn Gly Leu Ala Lys Glu Ser Glu Asp Asp Leu Asn Lys Glu Ser Glu	
	515 520 525	
45	GAG GAG GTT GGC CCC ACA AAA GAG TCC GAA GAA GAT GAC TCA GAG AAA	1692
	Glu Glu Val Gly Pro Thr Lys Glu Ser Glu Glu Asp Asp Ser Glu Lys	
	530 535 540 545	

	GAG TCT GAT GAA GAC TGC TCT GAA AAA CAG TCT GAA GAT GGC TCC GAA	1740
	Glu Ser Asp Glu Asp Cys Ser Glu Lys Gln Ser Glu Asp Gly Ser Glu	
	550 555 560	
5	AGA GAA TTT GAA GAA AAT GGT CTC GAG AAA GAT TTG GAC GAG GAA GGT	1788
	Arg Glu Phe Glu Glu Asn Gly Leu Glu Lys Asp Leu Asp Glu Glu Gly	
	565 570 575	
10	TCT GAA AAG GAG CTT CAT GAA AAT GTT CTT GAC AAA GAG TTA GAA GAA	1836
	Ser Glu Lys Glu Leu His Glu Asn Val Leu Asp Lys Glu Leu Glu Glu	
	580 585 590	
15	AAT GAC TCT GAA AAC TCC GAA TTT GAA GAT GAC GGC TCT GAA AAA GTG	1884
	Asn Asp Ser Glu Asn Ser Glu Phe Glu Asp Asp Gly Ser Glu Lys Val	
	595 600 605	
20	TTA GAT GAG GAA GGC TCT GAG AGA GAG TTT GAC GAA GAT TCA GAT GAA	1932
	Leu Asp Glu Glu Gly Ser Glu Arg Glu Phe Asp Glu Asp Ser Asp Glu	
	610 615 620 625	
25	AAG GAA GAA GAG GAG GAT ACA TAT GAA AAA GTA TTT GAT GAT GAG TCT	1980
	Lys Glu Glu Glu Glu Asp Thr Tyr Glu Lys Val Phe Asp Asp Glu Ser	
	630 635 640	
30	GAT GAG AAA GAG GAT GAA GAA TAT GCA GAT GAA AAG GGG CTT GAA GCT	2028
	Asp Glu Lys Glu Asp Glu Glu Tyr Ala Asp Glu Lys Gly Leu Glu Ala	
	645 650 655	
35	GCT GAT AAA AAG GCG GAA GAA GGT GAT GCA GAT GAA AAG CTG TTT GAA	2076
	Ala Asp Lys Lys Ala Glu Glu Gly Asp Ala Asp Glu Lys Leu Phe Glu	
	660 665 670	
40	GAG TCA GAT GAC AAG GAA GAT GAA GAT GCA GAT GGA AAG GAA GTT GAA	2124
	Glu Ser Asp Asp Lys Glu Asp Glu Asp Ala Asp Gly Lys Glu Val Glu	
	675 680 685	
45	GAT GCT GAC GAA AAG TTG TTC GAA GAT GAT GAT TCC AAT GAG AAG TTG	2172
	Asp Ala Asp Glu Lys Leu Phe Glu Asp Asp Asp Ser Asn Glu Lys Leu	
	690 695 700 705	
50	TTT GAT GAG GAG GAA GAT TCC AGT GAG AAG TTG TTT GAC GAT TCT GAT	2220
	Phe Asp Glu Glu Glu Asp Ser Ser Glu Lys Leu Phe Asp Asp Ser Asp	
	710 715 720	
55	GAG AGG GGG ACT TTG GGT GGT TTT GGG AGT GTT GAA GAA GGG CCC CTA	2268
	Glu Arg Gly Thr Leu Gly Gly Phe Gly Ser Val Glu Glu Gly Pro Leu	
	725 730 735	

TCC ACT GGC AGC AGC TTT ATT CTC AGT AGC GAT GAT GAT GAC GAT GAT 2316
Ser Thr Gly Ser Ser Phe Ile Leu Ser Ser Asp Asp Asp Asp Asp Asp
740 745 750

5 ATT TAATCCCTTA AACTTGCTTT TTAGGGAGAG TCCTCCATCT ACATTTGCCT GTGCTT 2375
Ile

CAGGGTAATT ACTAGTAGTG TTACATGAAC ATGTGCATAG TGGTAGGATG CCATCAGATT 2435
10 AAAGCATTGA AGTGTTCAT TGTTACCTGT ACCTAATGGT TTAAATATA TGTTAATTGA 2495
TTGTTTAGTT AAAATGTCAT AGTTACAATG CAAGTAACT GGATACTTGT TCTTTTGTCA 2555
GATTTGTTAA ATGCATGCAG AATAATATTT TTAAGAGTAT TGATTGAAGT TTGTGATATT 2615
CATCAATAAA AATGAGTTGA TAATATGCAG AAAGTAAAA AAAAAAAAAA AAAAAA 2672

CLAIMS

1. An isolated nucleic acid molecule encoding a Tat-Stimulatory Factor protein.
- 5 2. The isolated nucleic acid molecule of Claim 1
 - (a) which hybridizes under stringent conditions to a molecule consisting of the coding region of the nucleic acid sequence of SEQ ID NO:1 and which codes for the Tat-Stimulatory Factor protein
 - (b) nucleic acid molecules that differ from the nucleic acid molecules of (a) in
10 codon sequence due to the degeneracy of the genetic code, and
 - (c) complements of (a) and (b).
3. The isolated nucleic acid molecule of claim 1, wherein the isolated nucleic acid molecule comprises the coding region of SEQ ID NO:1.
- 15 4. The isolated nucleic acid molecule of claim 1, wherein the isolated nucleic acid molecule consists essentially of the coding region of SEQ ID NO:1.
5. An isolated nucleic acid molecule selected from the group consisting of (a) a unique
20 fragment of SEQ ID NO:1 between 12 and 2650 nucleotides in length, and (b) complements of "(a)".
6. The isolated nucleic acid molecule of claim 4, wherein the isolated nucleic acid molecule is a unique fragment of the coding region of SEQ. I.D. NO. 1 and wherein the isolated nucleic
25 acid molecule is selected from the group consisting of at least 14 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
7. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule
30 is selected from the group consisting of at least 15 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".

8. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule is selected from the group consisting of at least 16 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
- 5 9. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule is selected from the group consisting of at least 17 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
- 10 10. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule is selected from the group consisting of at least 18 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
11. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule is selected from the group consisting of at least 20 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
- 15 12. The isolated nucleic acid molecule of claim 6, wherein the isolated nucleic acid molecule is selected from the group consisting of at least 22 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
- 20 13. The isolated nucleic acid molecule of claim 5 or 6, wherein the isolated nucleic acid molecule is selected from the group consisting of between 12 and 32 contiguous nucleotides of (a) SEQ ID NO:1, and complements of "(a)".
- 25 14. A host cell transformed or transfected with an expression vector comprising the isolated nucleic acid molecule of claim 1, 2, 3 or 4, operably linked to a promoter.
15. An isolated polypeptide comprising a Tat-Stimulatory Factor protein or a functional fragment thereof.

30

16. The isolated polypeptide of claim 15 wherein the peptide is coded for by the isolated nucleic acid molecule of claim 1, 2, 3 or 4.
17. An isolated polypeptide that selectively binds a Tat-Stimulatory Factor protein.
- 5 18. The isolated polypeptide of claim 17 wherein the isolated polypeptide selectively binds a protein coded for by the isolated nucleic acid molecule of claim 1, 2, 3 or 4.
19. The isolated polypeptide of claim 18, wherein the isolated polypeptide is an Fab or F(ab)
10 fragment of an antibody.
20. The isolated polypeptide of claim 18, wherein the isolated polypeptide is a fragment of an antibody, the fragment including a CDR3 region selective for the protein.
- 15 21. The isolated polypeptide of claim 18, wherein the isolated polypeptide is a monoclonal antibody.
22. The isolated polypeptide of claim 18, wherein the isolated polypeptide is a kinase.
- 20 23. A method for influencing transcriptional activity in a cell comprising
contacting the cell with an agent that selectively binds to an isolated nucleic acid molecule of claim 1 or an expression product thereof, in an amount effective to influence transcriptional activity in said cell.
- 25 24. A method as claimed in claim 23, wherein the agent is a modified nucleic acid.
25. A method as claimed in claim 23, wherein the agent is a polypeptide.
26. A method for treating a subject infected with HIV comprising

administering to a subject in need of such treatment an agent that selectively binds to an isolated nucleic acid molecule of claim 1 or an expression product thereof, in an amount effective to inhibit transcriptional elongation of HIV-1 by Tat in the subject.

- 5 27. A method of screening for a compound which binds to Tat-SF1 comprising:
contacting Tat-SF1 with the compound, and
determining the binding of the compound to Tat-SF1.

28. The method of claim 27, wherein the compound is detectably labeled and the step of
10 determining the binding comprises detecting the detectably labeled compound bound to the
Tat-SF1.

29. The method of claim 27, wherein the step of determining the binding comprises detecting
a change in the biological activity of the Tat-SF1.

15

30. The method of claim 29, wherein the detecting comprises comparing Tat-SF1 alone and
with the compound in an assay of Tat-SF1 biological activity selected from the group consisting
of a Tat-SF1 mediated transcription assay, a Tat-SF1 immunoassay, and a Tat-SF1-TAR binding
assay.

20

31. The method of claim 27, wherein the compound is an oligonucleotide.

32. A method of screening for a compound which binds to Tat-SF1-associated kinase
comprising:

25

- contacting Tat-SF1 associated kinase with the compound, and
determining the binding of the compound to Tat-SF1-associated kinase.

33. The method of claim 32, wherein the compound is detectably labeled and the step of
determining the binding comprises detecting the detectably labeled compound bound to the
30 Tat-SF1-associated kinase.

34. The method of claim 32, wherein the step of determining the binding comprises detecting a change in the biological activity of the Tat-SF1-associated kinase.

35. The method of claim 34, wherein the detecting comprises comparing Tat-SF1 alone and
5 with the compound in an assay of Tat-SF1-associated kinase biological activity selected from the group consisting of a Tat-SF1 mediated transcription assay, a Tat-SF1-associated kinase immunoassay, and a Tat-SF1-associated kinase substrate phosphorylation assay.

36. The method of claim 32, wherein the compound is an oligonucleotide.

10

37. A method of screening for a compound which binds to a complex of Tat-SF1 and Tat-SF1-associated kinase comprising:

contacting the complex of Tat-SF1 and Tat-SF1-associated kinase with the compound,

and

15 determining the binding of the compound to the complex of Tat-SF1 and

Tat-SF1-associated kinase.

38. The method of claim 37, wherein the compound is detectably labeled and the step of determining the binding comprises detecting the detectably labeled compound bound to the
20 complex of Tat-SF1 and Tat-SF1-associated kinase.

39. The method of claim 37, wherein the step of determining the binding comprises detecting a change in the biological activity of the complex of Tat-SF1 and Tat-SF1 associated kinase.

25 40. The method of claim 39, wherein the detecting comprises comparing the complex of Tat-SF1 and Tat-SF1-associated kinase alone and with the compound in an assay of the biological activity of the complex of Tat-SF1 and Tat-SF1-associated kinase selected from the group consisting of a Tat-SF1 mediated transcription assay, an immunoassay of the complex of Tat-SF1 and Tat-SF1-associated kinase, a Tat-SF1-associated kinase substrate phosphorylation
30 assay, and a Tat-SF1-TAR binding assay.

41. The method of claim 37, wherein the compound is an oligonucleotide.

42. A method of screening for compounds which modulate Tat-SF1-mediated transcriptional activation, comprising:

- 5 providing a mammalian cell which comprises a gene encoding a Tat-SF1 polypeptide,
 contacting the cell with a compound, and
 determining Tat-SF1 mediated transcriptional activation as a measure of the modulation
of Tat-SF1 mediated transcriptional activation provided by the compound.

10 43. The method of claim 42, wherein the mammalian cell further comprises a gene encoding
a Tat polypeptide.

44. The method of claim 43, wherein the mammalian cell further comprises an indicator gene
encoding an indicator gene product, and wherein the indicator gene is operably linked to a TAR
15 element.

45. The method of claim 44, further comprising a gene encoding a Tat-SF1-associated kinase
polypeptide.

20 46. The method of claim 45, wherein the Tat-SF1 polypeptide, the Tat polypeptide, the
indicator gene product and the Tat-SF1-associated kinase polypeptide are encoded by expression
vectors, which expression vectors are transfected into the mammalian cell.

47. The method of claim 44, wherein the indicator gene consists essentially of a molecule
25 selected from the group consisting of a gene encoding β -galactosidase, a gene encoding alkaline
phosphatase, a gene encoding chloramphenicol acetyl transferase, a gene encoding luciferase,
and a gene encoding green fluorescent protein.

48. The method of claim 44, wherein the TAR element is an HIV-1 LTR TAR element.

30

1/9

Tat-SF: — +
Tat: — + — +

TAR-G400

ΔTAR-G100

1 2 3 4

Fig. 1A




Fig. 1B




Fig. 1C

SUBSTITUTE SHEET (RULE 28)





Fig. 2A

SUBSTITUTE SHEET (RULE 26)

3/9



SUBSTITUTE SHEET (RULE 26)





Fig. 3A

SUBSTITUTE SHEET (RULE 26)

5/9



GST-Tat column
Tat column




Fig. 3C

SUBSTITUTE SHEET (RULE 26)




Fig. 4A

pre-immune 2X
 α-pp140 Ab 2X




Fig. 4C

SUBSTITUTE SHEET (RULE 26)




Fig. 4B

SUBSTITUTE SHEET (RULE 26)

8/9




Fig. 5A

Tat-SF1: 30 TQTDAGGEPDSLGOQ 44
EWS: 209 SQQNTYGPSSYGOQ 223

Tat-SF1: 82 ASSSTANVEDVHARTAEFPQEKAPPTDARKKGEKRAES 122
EWS: 113 AAQSAYGTQPAYPAYGOQPAATAPTRPDGNKPTTTSQPQS 153

Tat-SF1: 128 EEDRNTNVVVSGLPPDITVDEFIQLMKFGIIMRDPQTEEFVKLYKDNQ 177
EWS: 356 EDSNSATVQGLNDSVTLDDLADFFKQCGVVKMKRTGQPMIHYLDKET 406

Tat-SF1: 178 GNLKRGDGLCCYLKRESVELALKLLDEDEIRGYKLHVEVAK 217
EWS: 407 GKPKGDATVSYEDPPTAKAAVEWFDGKDFQGSKLKVSILAR 446

Tat-SF1: 311 PDGVASVSFRDPEEADYCIQTL DGRWFGGRQI 342
EWS: 409 PKGDATVSYEDPPTAKAAVEWFDGKDFQGSKL 440

Fig. 5B

SUBSTITUTE SHEET (RULE 26)

9/9




Fig. 6A

Tat:	VECTOR			Tat-SFI			FOLD * ENHANCEMENT
	- [†]	+	FOLD ACT.	-	+	FOLD ACT.	
EXP. 1	100	7228	72.3	31.7	7699	242.9	3.36
EXP. 2	100	15779	157.8	17.0	16548	973.4	6.17
EXP. 3	100	4899	49.0	35.3	10353	293.3	5.99
TFEB-VP16:	-	+	FOLD ACT.	-	+	FOLD ACT.	
EXP. 1	100	30229	302.3	118	20782	176.1	0.58
EXP. 2	100	18241	182.4	179	15080	84.2	0.46
GAL4-VP16:	-	+	FOLD ACT.	-	+	FOLD ACT.	
	100	132208	1322	95.0	129960	1368	1.03

Fig. 6B

SUBSTITUTE SHEET (RULE 26)